

CROCIN MAX- acetaminophen capsule, gelatin coated
OPMX LLC

Crocin MAX Pain Relief

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever / Fever Reducer

Uses

- Temporarily relieves minor aches and pains due to:
- Minor pain of arthritis
- Muscular aches
- Backache
- Premenstrual and menstrual cramps
- The common cold
- Headache
- Toothache
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- More than 6 caplets in 24 hours, which is the maximum daily amount
- With other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- Skin reddening
- Blisters
- Rash

If skin reaction occurs, stop use and seek medical help right away.

Do not use

- With any other containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- Pain gets worse or lasts more than 10 days
- Fever gets worse or lasts more than 3 days
- New symptoms occur
- Redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out reach of children.

Keep out reach of children.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **Do not take more than directed (see overdose warning)**

Adults	Take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor
Under 18 years of age	Ask a doctor

Other information

- Store between 20-25° C (68-77° F)
- Do not use if carton is opened.
- Do not use if foil inner seal is broken or missing

Inactive ingredients

caranuba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose,

pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin.

Questions or comments?

Call **(619)600-5632** (Mon - Fri 9am - 5pm PST)

Made for:

RRX

San Diego, CA 92154

Exclusively distributed by:

OPMX

Chula Vista, CA 91910

Phone: 619-600-5632

Package Label

Crocin MAX

Acetaminophen 650 mg

Pain Relief

Alivia el Dolor / Pain Relief

Reduce la fiebre

Fever Reducer

Dolor de cabeza

Headache

Dolores Musculares

Muscle Aches

Dolor leve por Artritis

Minor Pain of Arthritis

24 Caplets

NO VANISH AREA

Drug Facts (continued)
 Stop use and ask a doctor if:
 ■ Pain gets worse or lasts more than 10 days
 ■ Fever gets worse or lasts more than 3 days
 ■ New symptoms occur
 ■ Redness or swelling is present
 These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.
 Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222)
 Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
 ■ Do not take more than directed (see overdose warning)
 Adults
 Take 2 caplets every 8 hours with water
 swallow whole; do not crush, chew, split or dissolve
 do not take more than 6 caplets in 24 hours
 Under 18 years
 Ask a doctor
 of age
Other Information
 ■ Store between 20-25°C (68-77°F)
 ■ Do not use if carton is opened.
 ■ Do not use if foil inner seal is broken or missing.
Inactive ingredients caranuba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, polydioxane powder, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin.
Questions or comments?
 Call (619)600-5632 (Mon - Fri 9am - 5pm PST)

Drug Facts
Active ingredient (in each caplet) Acetaminophen 650 mg
Purpose Pain reliever / Fever Reducer
Uses
 ■ Temporarily relieves minor aches and pains due to:
 ■ Minor pain of arthritis
 ■ Muscular aches
 ■ Backache
 ■ The common cold
 ■ Premenstrual and menstrual cramps
 ■ Headache
 ■ Toothache
 ■ Temporarily reduces fever
Warnings: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ More than 6 caplets in 24 hours, which is the maximum daily amount
 ■ With other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product.
Alergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ Skin redness ■ Blisters ■ Rash
 If skin reaction occurs, stop use and seek medical help right away.
Do not use
 ■ With any other containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ If you are allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if you have liver disease
 ■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Crocin
MAX
 Acetaminophen 650 mg

Pain Relief

NDC 69729-801-24

Crocin
MAX
 Acetaminophen 650 mg



Pain Relief / Alivia el Dolor

Fever Reducer
 Reduce la fiebre
Headache
 Dolor de Cabeza

Muscles Aches
 Dolores Musculares
Minor Pain or Arthritis
 Dolor leve por Artritis

24 Caplets

Crocin
MAX
 Acetaminophen 650 mg
Pain Relief

Lot No:
 Exp. Date:

NO VANISH AREA

TAMPER EVIDENT:
 Do not use if package is opened or if blister unit is torn, broken or shows any sign of tampering

REV 12.2025 ITEM RRX1158



Made for:
RRX
 San Diego, CA 92154

Exclusively distributed by:
PMX
 Chula Vista, CA 91910
 Phone: 619-600-5632

<p>Datos del medicamento</p> <p>Ingrediente activo (en cada cápsula) Propósito Acetaminofén 650 mg Alivio del dolor / Reductor de fiebre</p> <p>Usos</p> <ul style="list-style-type: none"> ■ Alivia temporalmente dolores y molestias menores debido a: ■ Dolor leve por artritis ■ Dolores musculares ■ Dolor de espalda ■ Cólicos premenstruales y menstruales ■ Resfriado común ■ Dolor de cabeza ■ Dolor de muelas ■ Reduce temporalmente la fiebre <p>Advertencias</p> <p>Advertencia sobre el hígado: Este producto contiene acetaminofén. Puede ocurrir daño hepático severo si usted toma:</p> <ul style="list-style-type: none"> ■ Más de 6 cápsulas en 24 horas, que es la cantidad máxima diaria ■ Con otros medicamentos que contengan acetaminofén ■ 3 o más bebidas alcohólicas cada día mientras usa este producto <p>Alerta de alergia: El acetaminofén puede causar reacciones severas en la piel. Los síntomas pueden incluir:</p> <ul style="list-style-type: none"> ■ Enrojecimiento de la piel ■ Ampollas ■ Erupción cutánea <p>Si ocurre una reacción en la piel, suspenda su uso y busque ayuda médica de inmediato.</p> <p>No usar</p> <ul style="list-style-type: none"> ■ Con ningún otro producto que contenga acetaminofén (de prescripción o sin receta). Si no está seguro de si un medicamento contiene acetaminofén, consulte a un médico o farmacéutico. ■ Si es alérgico al acetaminofén o a alguno de los ingredientes inactivos en este producto <p>Consulte a un médico antes de usar si tiene enfermedad hepática</p> <p>Consulte a un médico o farmacéutico antes de usar si está tomando el medicamento anticoagulante warfarina</p>	<p>Datos del medicamento (continuación)</p> <p>Suspenda el uso y consulte a un médico si</p> <ul style="list-style-type: none"> ■ El dolor empeora o dura más de 10 días ■ La fiebre empeora o dura más de 3 días ■ Aparecen nuevos síntomas ■ Hay enrojecimiento o hinchazón <p>Estos podrían ser signos de una condición seria.</p> <p>Si está embarazada o amamantando, consulte a un profesional de salud antes de usar. Mantener fuera del alcance de los niños.</p> <p>Advertencia de sobredosis: en caso de sobredosis, obtenga ayuda médica o contacte a un Centro de Control de Envenenamientos de inmediato (1-800-222-1222). La atención médica rápida es crucial tanto para adultos como para niños, incluso si no nota ningún signo o síntoma.</p> <p>Instrucciones</p> <ul style="list-style-type: none"> ■ No tome más de lo indicado (ver advertencia de sobredosis) <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20%;">Adultos</td> <td>Tome 2 cápsulas cada 8 horas con agua tráguelas enteras; no las triture, mastique, parta ni disuelva no tome más de 6 cápsulas en 24 horas no use por más de 10 días a menos que lo indique un médico</td> </tr> <tr> <td>Menores de 18 años</td> <td>Consulte a un médico</td> </tr> </table> <p>Otra información</p> <ul style="list-style-type: none"> ■ Almacene entre 20-25°C (68-77°F) ■ No use si el empaque está abierto. ■ No usar si el sello interior de aluminio está roto o ausente. <p>Ingredientes inactivos Cera de carnauba, hidroxietil celulosa, hipromelosa, estearato de magnesio, celulosa microcristalina, almidón modificado, povidona, celulosa pulverizada, almidón pregelatinizado, glicolato de almidón sódico, dióxido de titanio, triacetina.</p> <p>¿Preguntas o comentarios? Llame al (619)600-5632 (Mon- Fri 9am - 5pm PST)</p>	Adultos	Tome 2 cápsulas cada 8 horas con agua tráguelas enteras; no las triture, mastique, parta ni disuelva no tome más de 6 cápsulas en 24 horas no use por más de 10 días a menos que lo indique un médico	Menores de 18 años	Consulte a un médico
Adultos	Tome 2 cápsulas cada 8 horas con agua tráguelas enteras; no las triture, mastique, parta ni disuelva no tome más de 6 cápsulas en 24 horas no use por más de 10 días a menos que lo indique un médico				
Menores de 18 años	Consulte a un médico				

CROCIN MAX

acetaminophen capsule, gelatin coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-801
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
TRIACETIN (UNII: XHX3C3X673)	
HYDROXYMETHYL CELLULOSE (UNII: 273FM27VK1)	
POVIDONE K30 (UNII: U725QWY32X)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL (Oblong shape)	Size	21mm
Flavor		Imprint Code	OPMX
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-801-24	2 in 1 CARTON	01/29/2026	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/29/2026	

Labeler - OPMX LLC (029918743)

Establishment

Name	Address	ID/FEI	Business Operations
HBC HEALTHCARE PRIVATE LIMITED		854255181	manufacture(69729-801)

Revised: 1/2026

OPMX LLC