

**MEJORALITO CHILDREN CHILDREN- acetaminophen tablet, chewable  
OPMX LLC**

-----  
**Mejoralito chewable CHILDREN**

***Drug Facts***

***Active ingredients (in each chewable tablet)***

Acetaminophen 160 mg

***Purposes***

Pain reliever/Fever reducer

***Uses***

Temporarily relieves minor aches and pains due to:

- Headache
- Toothache
- Flu
- Sore throat
- The common cold
- Temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if your child takes.

- More than 5 doses in 24 hours, which is the maximum daily amount
- With other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

***Do not use***

With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist.

If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin.

**When using this product do not exceed recommended dosage.**

Ask a doctor before use if your child has liver disease.

**Stop use and ask a doctor if**

- Pain gets worse or lasts more than 5 days
- Fever gets worse or lasts more than 3 days
- New symptoms occur
- Redness or swelling is present.

These could be signs of a serious condition.

**Keep out of reach of children.**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **This product does not contain directions or complete warnings for adult use.**
- **Do not give more than directed.**
- Find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- Chew or crush tablets completely before swallowing; do not swallow tablets whole.
- Repeat dose every 4 hours while symptoms last.
- Do not give more than 5 times in 24 hours.
- Do not give for more than 5 days unless directed by a doctor.

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

\*or as directed by a doctor

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- Avoid high humidity
- Store at 25°C (77°F); excursions permitted between 15° -30°C (59° -86°F)
- See and flap for expiration date and lot number

**Inactive Ingredients**

Citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminium lake, dextrans hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose.

**Questions or comments?**

Call **(619)600-5632** (Mon- Fri 9am - 5pm PST)

Exclusively distributed by:

OPMX

Chula Vista, CA 91910

**Phone: 619-600-5632**

inov

FARMACEUTIQUE

**PRINCIPAL DISPLAY PANEL**

NDC 69729-160-24

Mejoralito®

CHILDREN

Acetaminophen 160 mg

Auxiliar en el alivio del  
DOLOR y FIEBRE

Pain reliever/

Fever Reducer

- Sin Aspirina / Aspirin Free
- Sin Ibuprofeno / Ibuprofen Free

2-11 AÑOS/YEARS

24 Tabletas Masticables

Chewable Tablets

Sabor Uva/Grape Flavor

Acetaminophen 160 mg  
Chevable Tablets

Pain Reliever /  
Fever Reducer

**Mejoralito**  
CHILDREN

**Mejoralito**  
CHILDREN

NDC 69729-160-24

**Acetaminophen 160 mg**

**Auxiliar en el alivio del  
DOLOR y FIEBRE**

Pain Reliever /  
Fever Reducer

- Sin Aspirina / Aspirin Free
- Sin Ibuprofeno / Ibuprofen Free

**2-11**  
AÑOS/YEARS

24 **Tabletas Masticables**  
Chewable Tablets

Sabor Uva / Grape Flavor

Acetaminophen 160 mg  
Chewable Tablets

**Mejoralito**  
CHILDREN

Batch:  
Exp Date:

NO VARNISH AREA

Exclusively distributed by:






Rev. 05.2025 Item #: IN0341

Acetaminophen 160 mg  
Chewable Tablets

**Mejoralito**  
CHILDREN

NO VARNISH AREA

**Drug Facts**

**Active Ingredients**      **Purposes**  
(in each chewable tablet)

Acetaminophen 160 mg..... Pain reliever/  
Fever reducer

**Uses** Temporarily relieves minor aches and pains due to:

- Headache
- Sore throat
- Toothache
- The common cold
- Flu
- Temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- More than 5 doses in 24 hours, which is the maximum daily amount.
- With other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product do not exceed recommended dosage.** Ask a doctor before use if your child has liver disease.

**Stop use and ask a doctor if**

- Pain gets worse or lasts more than 5 days
- Fever gets worse or lasts more than 3 days
- New symptoms occur
- Redness or swelling is present.

These could be signs of a serious condition.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- This product does not contain directions or complete warnings for adult use.
- Do not give more than directed.
- Find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- Chew or crush tablets completely before swallowing; do not swallow tablets whole.
- Repeat dose every 4 hours while symptoms last.
- Do not give more than 5 times in 24 hours.
- Do not give for more than 5 days unless directed by a doctor.

**Drug Facts (continued)**

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

\*or as directed by a doctor

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- Avoid high humidity
- Store at 25°C (77°F); excursions permitted between 15° -30°C (59°-86°F)
- See and flap for expiration date and lot number

**Inactive Ingredients**

Citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose.

**Questions or comments?**  
Call (619)600-5632 (Mon- Fri 9am - 5pm PST)

NO VARNISH AREA

NO ABRIR AQUÍ

**Datos del medicamento**

**Ingredientes Activos (in each chewable tablet)** **Propósito**

Acetaminofén 160 mg ..... Analgésico/  
Reductor de Fiebre

**Usos** Alivia temporalmente dolores menores y molestias debido a:

- Dolor de Cabeza
- Dolor de Muela
- Flu
- Dolor de Garganta
- Resfriado Común
- Reduce temporalmente la fiebre

**Advertencias**

**Advertencia sobre el hígado:** Este producto contiene acetaminofén. Puede causar daño hepático grave si tu hijo toma:

- Más de 5 dosis en 24 horas (límite máximo diario)
- Otros medicamentos que también contengan acetaminofén

**Advertencia de alergia:** El acetaminofén puede causar reacciones cutáneas graves. Los síntomas pueden incluir:

- Enrojecimiento de la piel
- Ampollas
- Erupción cutánea

Si ocurre una reacción en la piel, suspender el uso y buscar ayuda médica de inmediato.

**Advertencia sobre el dolor de garganta:** Si el dolor de garganta es severo, dura más de 2 días o se acompaña de fiebre, dolor de cabeza, sarpullido, náuseas o vómitos, consultar a un médico.

**No usar**

Con otros medicamentos que contengan acetaminofén (de prescripción o sin receta). Si no estás seguro, consulta con un médico o farmacéutico.

Si tu hijo es alérgico al acetaminofén o a alguno de los ingredientes inactivos de este producto.

**Datos del medicamento (continuación)**

Consulta a un médico o farmacéutico antes de usar este medicamento si tu hijo está tomando el anticoagulante warfarina.

**Cuando uses este producto, no excedas la dosis recomendada.** Consulta a un médico antes de usar este medicamento si tu hijo tiene enfermedad hepática.

**Suspenda el uso y consulte a un médico si:**

- El dolor empeora o dura más de 5 días.
- La fiebre empeora o dura más de 3 días.
- Aparecen nuevos síntomas.
- Hay enrojecimiento o hinchazón.

Estos podrían ser signos de una condición grave.

**Manténgase fuera del alcance de los niños.** En caso de sobredosis, busque ayuda médica o comuníquese de inmediato con un Centro de Control de Envenenamientos. La atención médica rápida es crucial tanto para adultos como para niños, incluso si no nota ningún signo o síntoma.

**Indicaciones**

- Este producto no contiene instrucciones ni advertencias completas para el uso en adultos.
- No dé más de lo indicado.
- Encuentre la dosis correcta en la tabla a continuación. Si es posible, utilice el peso para la dosis; de lo contrario, utilice la edad.
- Mastique o triture completamente las tabletas antes de tragarlas; no trague las tabletas enteras.
- Repita la dosis cada 4 horas mientras duren los síntomas.
- No dé más de 5 veces en 24 horas.
- No dé por más de 5 días a menos que lo indique un médico.

**Datos del medicamento (continuación)**

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Menos de 24	Menos de 2	Consulte a un doctor
24-35	2-3	1 tableta
36-47	4-5	1 1/2 tabletas
48-59	6-8	2 tabletas
60-71	9-10	2 1/2 tabletas
72-95	11	3 tabletas

\*o según lo indicado por un médico

**Otra Información**

- **EVIDENCIA DE ALTERACIÓN: NO USE SI EL PAQUETE EXTERIOR ESTÁ ABIERTO O EL BLISTER ESTÁ ROTO O QUEBRADO.**
- Evite la alta humedad.
- Conservar a 25°C (77°F); se permiten variaciones entre 15° -30°C (59°-86°F).
- Consulte el reverso para la fecha de vencimiento y el número de lote.

**Ingredientes Inactivos**

Ácido cítrico, crospovidona, D&C rojo #27 laca de aluminio, D&C rojo #30 laca de aluminio, dextratos hidratados, etilcelulosa, FD&C azul #1 laca de aluminio, sabores, estearato de magnesio, manitol, polietileno, ácido estéarico, sacarosa.

**¿Preguntas o comentarios?**  
Llame al (619)600-5632 (Mon- Fri 9am - 5pm PST)

NO ABRIR AQUÍ

## MEJORALITO CHILDREN CHILDREN

acetaminophen tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69729-160
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>DEXTRATES</b> (UNII: G263MI44RU)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>D&amp;C RED NO. 30 ALUMINUM LAKE</b> (UNII: GE75M6YV5W)	
<b>ETHYLCELLULOSE</b> (UNII: 7Z8S9VYZ4B)	
<b>CITRIC ACID</b> (UNII: 2968PHW8QP)	
<b>D&amp;C RED NO. 27 ALUMINUM LAKE</b> (UNII: ZK64F7XSTX)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>POLYETHYLENE</b> (UNII: UG00KM4WR7)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3Z D4)	

## Product Characteristics

<b>Color</b>	purple (Light purple/pink)	<b>Score</b>	no score
<b>Shape</b>	OVAL (Oblong)	<b>Size</b>	17mm
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	OPMX
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-160-24	2 in 1 CARTON	01/30/2026	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/30/2026	

**Labeler** - OPMX LLC (029918743)

## Establishment

Name	Address	ID/FEI	Business Operations
HBC HEALTHCARE PRIVATE LIMITED		854255181	manufacture(69729-160)

Revised: 1/2026

OPMX LLC