

RELAXINK LIDOCAINE- lidocaine 4% cream
Hambli Limited

Warnings

(For external use only)

When using this product:

Avoid contact with eyes.

Do not exceed recommended dosage unless directed by a doctor.

Do not put this product into the rectum by using fingers or any mechanical device or applicator.

Stop use and ask a doctor if:

Rectal bleeding occurs.

Condition worsens or does not improve within 7 days.

Allergic reaction occurs to ingredients in this product.

Symptom being treated does not subside or if redness, irritation, swelling, pain or other symptoms develop or increase symptoms clear up and return within a few days.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

When practical, clean area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or soft cloth before applying.

Adults and children 12 years and older:

apply externally to the affected area up to 6 times a day.

For the temporary relief of pain and discomfort associated with anorectal disorders.

Keep out of reach of children.

Inactive Ingredients

Water, Glycerin, Propylene Glycol, Hydroxyethyl Acrylate/
Sodium Acryloyldimethyl Taurate Copolymer, Caprylic/
Capric Triglyceride, Tocopheryl Acetate, Arginine,
Phenoxyethanol, Chlorphenesin

Uses

Helps relieve the pain, itching and burning associated with hemorrhoids and other anorectal disorders.

Avoid contact with eyes.

Do not exceed recommended dosage unless directed by a doctor.

Do not put this product into the rectum by using fingers or any mechanical device or applicator.

Stop use and ask a doctor if:

Rectal bleeding occurs.
Condition worsens or does not improve within 7 days.
Allergic reaction occurs to ingredients in this product.
Symptom being treated does not subside or if redness irritation, swelling, pain or other symptoms develop or increase symptoms clear up and return within a few days.
If pregnant or breastfeeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Local anesthetic

Lidocaine 4%.

Pregnancy Section

Consult your doctor before using it

DRUG FACTS

DRUG FACTS

Active Ingredient

Lidocaine 4%.

Uses

Helps relieve the pain, itching and burning associated with hemorrhoids and other anorectal disorders.

Warnings (For external use only)

When using this product:

Avoid contact with eyes.

Do not exceed recommended dosage unless directed by a doctor.

Do not put this product into the rectum by using fingers or any mechanical device or applicator.

Stop use and ask a doctor if:

Rectal bleeding occurs.

Condition worsens or does not improve within 7 days.

Allergic reaction occurs to ingredients in this product.

Symptom being treated does not subside or if redness irritation, swelling, pain or other symptoms develop or increase symptoms clear up and return within a few days.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Purpose

Local anesthetic

DRUG FACTS (Continued)

Directions

When practical, clean area with mild soap and warm

water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or soft cloth before applying.
Adults and children 12 years and older:
apply externally to the affected area up to 6 times a day.
Children under 12 years of age: Consult a doctor.
To use finger cots: Roll one finger cot over finger.
Gently squeeze cream onto finger cot. Smooth a layer of the cream over affected area.

Other Information

Store at 25°C (77°F); excursions permitted to 15-30C (59-86°F) [see USP Controlled Room Temperature].

Inactive Ingredients

Water, Glycerin, Propylene Glycol, Hydroxyethyl Acrylate/
Sodium Acryloyldimethyl Taurate Copolymer, Caprylic/
Capric Triglyceride, Tocopheryl Acetate, Arginine,
Phenoxyethanol, Chlorphenesin



DRUG FACTS

Active Ingredient	Purpose
Lidocaine 4%	Local anesthetic

Uses

- Helps relieve the pain, itching and burning associated with hemorrhoids and other anorectal disorders.

Warnings (For external use only)

When using this product:

- Avoid contact with eyes.
- Do not exceed recommended dosage unless directed by a doctor.
- Do not put this product into the rectum by using fingers or any mechanical device or applicator.

Stop use and ask a doctor if:

- Rectal bleeding occurs.
- Condition worsens or does not improve within 7 days.
- Allergic reaction occurs to ingredients in this product.
- Symptom being treated does not subside or if redness, irritation, swelling, pain or other symptoms develop or increase symptoms clear up and return within a few days.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

DRUG FACTS (Continued)

Directions

- When practical, clean area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or soft cloth before applying.
- **Adults and children 12 years and older:** apply externally to the affected area up to 6 times a day.
- **Children under 12 years of age:** Consult a doctor.
- **To use finger cots:** Roll one finger cot over finger. Gently squeeze cream onto finger cot. Smooth a layer of the cream over affected area.

Other Information

- Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

Inactive Ingredients

Water, Glycerin, Propylene Glycol, Hydroxyethyl Acrylate/Sodium Acryloyldimethyl Taurate Copolymer, Caprylic/Capric Triglyceride, Tocopheryl Acetate, Arginine, Phenoxyethanol, Chlorphenesin



RELAXINK LIDOCAINE

lidocaine 4% cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83714-0004
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
CHLORPHENESIN (UNII: I670DAL4SZ)	0.15 g in 100 g
GLYCERIN (UNII: PDC6A3C0OX)	7 g in 100 g
CAPRYLIC/CAPRIC/LAURIC TRIGLYCERIDE (UNII: FJ1H6M2JG9)	2 g in 100 g
PHENOXYETHANOL (UNII: HIE492ZZ3T)	0.6 g in 100 g
ARGININE (UNII: 94ZLA3W45F)	0.7 g in 100 g
.ALPHA.-TOCOPHEROL ACETATE, D- (UNII: A7E6112E4N)	1 g in 100 g
WATER (UNII: 059QF0KO0R)	76.35 g in 100 g
HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER (45000 MPA.S AT 1%) (UNII: 86FQE96TZ4)	3.2 g in 100 g
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	5 g in 100 g

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83714-0004-1	1 in 1 PACKAGE	02/15/2026	
1	NDC:83714-0004-0	30 g in 1 TUBE; Type 1: Convenience Kit of Co-Package		



DRUG FACTS		DRUG FACTS (Continued)	
Active Ingredient Lidocaine 4%	Purpose Local anesthetic	Directions	
Uses ■ Helps relieve the pain, itching and burning associated with hemorrhoids and other anorectal disorders.		<ul style="list-style-type: none"> ■ When practical, clean area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or soft cloth before applying. ■ Adults and children 12 years and older: apply externally to the affected area up to 6 times a day. ■ Children under 12 years of age: Consult a doctor. ■ To use finger cots: Roll one finger cot over finger. Gently squeeze cream onto finger cot. Smooth a layer of the cream over affected areas. 	
Warnings (For external use only) <i>When using this product:</i>		Other Information	
<ul style="list-style-type: none"> ■ Avoid contact with eyes. ■ Do not exceed recommended dosage unless directed by a doctor. ■ Do not put this product into the rectum by using fingers or any mechanical device or applicator. 		<ul style="list-style-type: none"> ■ Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. 	
Stop use and ask a doctor if:		Inactive Ingredients	
<ul style="list-style-type: none"> ■ Rectal bleeding occurs. ■ Condition worsens or does not improve within 7 days. ■ Allergic reaction occurs to ingredients in this product. ■ Symptom being treated does not subside or if redness, irritation, swelling, pain or other symptoms develop or increase symptoms clear up and return within a few days. 		Water, Glycerin, Propylene Glycol, Hydroxyethyl Acrylate/Sodium Acryloyldimethyl Taurate Copolymer, Caprylic/Capric Triglyceride, Tocopheryl Acetate, Arginine, Phenoxyethanol, Chlorphenesin	
<i>If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</i>			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/15/2026	

Labeler - Hambli Limited (227713374)

Establishment

Name	Address	ID/FEI	Business Operations
NANTONG HEALTH & BEYOND HYGIENIC PRODUCTS INC.		421280161	manufacture(83714-0004)

