

PRO NUMB SENSITIVE SKIN- numbing spray spray

Pro Numb Tattoo Numbing Spray LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Pro Numb Tattoo Sensitive Skin Spray 120 mL

Active Ingredient(s)

Lidocaine HCl 4.2% Epinephrine HCl .01%

Purpose

Topical Anesthetic
Vasoconstriction

Use

Temporarily takes away pain, itching, and swelling associated with anorectal disorders or pain-sensitive procedures.

Warnings

External Use Only. Keep out of reach of children.

Do not use if you have • seizures or liver disease • if pregnant or breastfeeding • a known allergy or sensitivity occurs, discontinue use, and seek medical attention. If the condition intensifies or symptoms persist for more than seven days or clear up and occur again within a few days, Cease the use of this product and consult a doctor. Do not use in large quantities, Particularly over raw surfaces or blistered areas. Avoid contact with the eyes. If contact should occur, irrigate with large amounts of sterile saline and seek medical attention. Do not swallow, if swallowed, seek medical attention immediately.

When using this product, you may notice temporary whitening or redness of the skin where Pro Numb is applied.

Keep out of reach of children.

Directions

A sensitivity test is encouraged before using. Spray a small amount on the broken area of the skin. Wait 60 seconds for the anesthetic to start working. Wash Pro Numb off completely before tattooing.

Other information


Store in a cool dark place or refrigerate and keep out of reach of children. Discard the item after the expiration date is due.

Inactive ingredients

Purified RO water, Paraben-DU, Disodium EDTA, Citric acid, Optiphen.

Contact Pro Numb Distribution. (800) 865-4965

Package Label - Principal Display Panel



Drug Fact- For Use by Licensed Tattoo artists only	
Active Ingredients	
Lidocaine HCl	4.2%
Epinephrine HCl	.01%
Purpose	
Topical Anesthetic	
Vasoconstriction	
Uses: External Use Only.	
Temporarily takes away pain, itching, and swelling associated with anorectal disorders or pain-sensitive procedures.	
Warnings-Keep out of reach of children	
Do not use if you have	
<ul style="list-style-type: none">• seizures or liver disease• if pregnant or breastfeeding• a known allergy or sensitivity occurs, discontinue use, and seek medical attention. If the condition intensifies or symptoms persist for more than seven days or clear up and occur again within a few days, Cease the use of this product and consult a doctor. Do not use in large quantities, Particularly over raw surfaces or blistered areas.	
Avoid contact with the eyes. If contact should occur, irrigate with large amounts of sterile saline and seek medical attention.	
DO NOT SWALLOW. If swallowed, seek medical attention immediately.	
When using this product you may notice temporary whitening or redness of the skin where pro numb is applied.	
Directions: Sensitivity test encouraged before use. spray a small amount on the broken area of the skin. Wait 60 seconds for the anesthetic to start working. Wash Pro Numb off completely before tattooing.	
Inactive Ingredients: purified RO water, Paraben-DU, Disodium EDTA, Citric acid, Optiphen.	
Other information: Store in a cool dark place or refrigerate and keep out of reach of children. Discard after the expiration date.	
Questions? Contact pro numb Distribution	
Distributed by: Pro Numb Tattoo numbing spray LLC Melbourne, FL, 32907 (800) 865-4965 www.tattoonumbingsprays.com	

PRO NUMB SENSITIVE SKIN

numbing spray spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83389-112
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EPINEPHRINE HYDROCHLORIDE (UNII: WBB0470038) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE HYDROCHLORIDE	0.1 mg in 100 mL
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	4.2 mg in 100 mL

Inactive Ingredients

Ingredient Name	Strength
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATER (UNII: 059QF0KO0R)	
EDETA TE DISODIUM (UNII: 7FLD91C86K)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83389-112-11	120 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	02/02/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		02/02/2026	

Labeler - Pro Numb Tattoo Numbing Spray LLC (118964052)

Revised: 2/2026

Pro Numb Tattoo Numbing Spray LLC