#### PAIN RELIEVER PM EXTRA STRENGTH- acetaminophen and diphenhydramine hydrochloride tablet, coated Marc Glassman, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### 1095 - MAR - 2018-1206

## **Drug Facts**

Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever
Diphenhydramine HCl 25 mg	Nighttime sleep aid

#### Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

## Warnings

#### Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

## Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

## Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

## If pregnant or breast-feeding, ask a health professional before use.

## Keep out of reach of children.

## **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

## • do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul> <li>take 2 caplets at bedtime</li> <li>do not take more than 2 caplets of this product in 24 hours</li> </ul>
children under 12 years	<ul> <li>do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage</li> </ul>

## Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

## Inactive ingredients

colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1,

hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

## PRINCIPAL DISPLAY PANEL

†Compare to the active ingredient in Tylenol® PM Extra Strength Caplets

Marc's®

Extra Strength


Pain Reliever PM

Pain Reliever/Nighttime Sleep-Aid

Acetaminophen, Diphenhydramine HCI

- Contains No Aspirin
- Non-Habit Forming

50 Caplets



# PAIN RELIEVER PM EXTRA STRENGTH

acetaminophen and diphenhydramine hydrochloride tablet, coated

<b>Product Information</b>					
Product Type	HUMAN OTC DRUG	Item Code (	Source)	NDC:6899	8-095
Route of Administration	ORAL				
	<b>14</b> - <b>1</b> - <b>1</b>				
Active Ingredient/Active	Molety				
Ingre	dient Name		Basis of Str	ength	Strength
ACETAMINOPHEN (UNII: 36209IT	L9D) (ACETAMINOPHEN - UN	ll:36209ITL9D)	ACETAMINOPHEN		500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)		DIPHENHYDRAMIN	E	~-	
	583M)		HYDROCHLORIDE		25 mg

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
COPOVIDONE K25-31 (UNII: D9C330MD8B)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: 08232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

#### **Product Characteristics**

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AAA;1031
Contains			

## Packaging

# Item Code	Package Description	Marketing Start Date	Marketing End Date
<b>1</b> NDC:68998- 095-02	1 in 1 CARTON	03/01/2008	02/28/2025
1	50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing	Information		
Marketing Marketing Category	Information Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

Labeler - Marc Glassman, Inc. (094487477)

Revised: 11/2022

Marc Glassman, Inc.