

**INFANTS TYLENOL - acetaminophen suspension**  
**Kenvue Brands LLC**

-----  
**Infants' TYLENOL®**

Drug Facts

**Active ingredient**

**Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

**Purpose**

**Purpose**

Pain reliever/fever reducer

**Uses**

**Uses** temporarily:

- reduces fever
- relieves minor aches and pains due to:

-the common cold

-flu

-headache

-sore throat

-toothache

**Warnings**

**Warnings**

**Liver warning**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if your child has** liver disease

**Ask a doctor or pharmacist before use if your child is** taking the blood thinning drug warfarin

**When using this product do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**Directions**

- **this product does not contain directions or complete warnings for adult use.**

- **do not give more than directed (see overdose warning)**

- **shake well before using**

- mL = milliliter

- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

- push air out of syringe. Insert syringe tip into bottle opening

- flip bottle upside down. Pull yellow part of syringe to correct dose

- dispense liquid slowly into child's mouth, toward inner cheek

- repeat dose every 4 hours while symptoms last

- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

### Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL

\* or as directed by a doctor

**Attention:** use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

### Other information

#### Other information

- each 5 mL contains: **potassium 5 mg**
- store between 20-25°C (68-77°F)
- **do not use if carton is opened**
- **do not use if bottle wrap imprinted with “TYLENOL” is broken or missing**

### Inactive ingredients

anhydrous citric acid, flavors, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

**Questions or comments?** call **1-800-458-1635** (toll-free) or **215-273-8755** (collect)

### NEW

NDC 50580-154-02

### Infant's TYLENOL®

For children

### Acetaminophen (160 mg per 5 mL)

Oral Suspension

Pain Reliever-Fever Reducer

### Pain+Fever

Infants

### DYE-FREE

**Free Of:**

- Dyes
- Alcohol
- Ibuprofen
- Aspirin
- Parabens
- High Fructose Corn Syrup
- Artificial Flavoring

**Natural Apple  
Flavor with Other  
Natural Flavors**

2 fl oz (60 mL) 160 mg per 5 mL\*

\*Same as Children's TYLENOL®

**SimpleMeasure®**

Specially designed for younger children  
Use Only Enclosed Syringe



How can we help?  
1-800-458-1635

OPEN HERE

Scan QR code to learn more about TYLENOL®



Important: Read all product information before using. Keep this box for important information.

### Drug Facts

**Active ingredient (in each 5 mL)** **Purpose**  
Acetaminophen 160 mg Pain reliever/fever reducer

#### Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

#### Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- this product does not contain directions or complete warnings for adult use.
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter

### Drug Facts (continued)

- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening
- flip bottle upside down. Pull yellow part of syringe to correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

#### Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL

\* or as directed by a doctor

**Attention:** use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

#### Other information

- each 5 mL contains: potassium 5 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened
- do not use if bottle wrap imprinted with "TYLENOL" is broken or missing

#### Inactive Ingredients

anhydrous citric acid, flavors, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

#### Questions or comments?

call 1-800-458-1635 (toll-free) or 215-273-8755 (collect)



NDC 50580-154-02

NEW

# Infants' TYLENOL®

For Children

## Acetaminophen (160 mg per 5 mL)

Oral Suspension

Pain Reliever-Fever Reducer

### Pain + Fever Infants

#### DYE-FREE

Free Of:

- Dyes
- Alcohol
- Ibuprofen
- Aspirin
- Parabens
- High Fructose Corn Syrup
- Artificial Flavoring

#### Natural Apple Flavor with Other Natural Flavors



SimpleMeasure®  
Specially designed for younger children

Use Only Enclosed Syringe

2 fl oz (60 mL)  
160 mg per 5 mL\*

\*Same as Children's TYLENOL®

# Infants' TYLENOL®

For Children

Pediatricians recommend TYLENOL® for pain and fever

#### IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose

#### SimpleMeasure®

STEP 1. Push air out of syringe.



STEP 2. Insert syringe tip into bottle opening.



STEP 3. Flip bottle upside-down.



STEP 4. Pull yellow part of syringe to correct dose.



Distributed by:  
Kenvue Brands LLC,  
Summit, NJ 07901  
© Kenvue Brands LLC 2026

Visit us at [www.tylenol.com](http://www.tylenol.com) 30062483  
Pat. [www.kenvuepats.com](http://www.kenvuepats.com)



## INFANTS TYLENOL

acetaminophen suspension

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:50580-154

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	160 mg in 5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
SUCROSE (UNII: C151H8M554)				
XANTHAN GUM (UNII: TTV12P4NEE)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
GLYCERIN (UNII: PDC6A3C0OX)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				
WATER (UNII: 059QF0KOOR)				
Product Characteristics				
Color	white (White to off-white)	Score		
Shape		Size		
Flavor	APPLE	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-154-02	1 in 1 CARTON	04/13/2026	
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	04/13/2026		

**Labeler** - Kenvue Brands LLC (118772437)