

**ACETAMINOPHEN 325MG- acetaminophen tablet**  
**Marlex Pharmaceuticals Inc**

-----  
**Acetaminophen 325mg**

***Drug Facts***

***Active Ingredient(in each tablet)***

Acetaminophen USP 325 mg

***Purpose***

*Pain Reliever/Fever Reducer*

***Uses***

To reduce fever and for the temporary relief of minor aches and pains due to:

- Headache
- Backache
- The common cold
- Premenstrual and menstrual cramps
- Temporarily reduces fever
- Muscular ache
- Minor pain of arthritis
- Toothache

**Warnings**

**Liver warning:**this product contains acetaminophen.

Severe liver damage may occur if

- adult takes more than 12 tablets in 24 hours, which is the maximum daily amount
- child takes more than 5 doeses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a Doctor**

**before use if you have** liver disease.

**Ask a doctor or pharmacist before use**

**if you are** taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- Pain gets worse or lasts more than 10 days in adults and children.
- Pain gets worse or lasts more than 5 days in children under 12 years.
- Fever gets worse or lasts more than 3 days.
- New symptoms occur.
- Redness or swelling is present.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.****Overdose warning:**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions****Do not take more than directed****Adults and Children 12 years and over:**

- Take 2 tablets every 4 to 6 hours while symptoms last.
- Do not take more than 12 tablets in 24 hours

**Children 6-11 years:**

- Take 1 tablet every 4 to 6 hours while symptoms last.
- Do not take more than 5 tablets in 24 hours.

**Children under 6years:**

Do not use this adult regular strength product in children under 6 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and may cause liver damage.

**Other Information**

- Store between 20-25 °C (68-77 °F).

**Inactive Ingredients**

Pregelatinized Starch, Povidone K 30, Sodium starch Glycolate, Stearic Acid.

**Manufactured for/ Distributed by:**

Marlex Pharmaceuticals, Inc.  
New Castle, DE 19720

Rev. 01/26 AA

**PRINCIPAL DISPLAY PANEL**

NDC 10135-0828-01

**Acetaminophen**

**325 mg**

**100 TABLETS**



<b>Drug Facts</b>	<b>Active ingredient (in each tablet)</b> Acetaminophen USP 325 mg...Pain reliever/Fever Reducer	<b>Purpose</b> Pain relief and fever reduction.
	<b>Uses</b> To reduce fever and for the temporary relief of minor aches and pains due to: • Headache • Backache • The common cold • Premenstrual and menstrual cramps	
	• Muscular aches • Minor pain of arthritis • Toothache • Temporarily reduces fever	
	<b>Warnings</b> <b>Liver Warning:</b> this product contains acetaminophen. Severe liver damage may occur if: • adult takes more than 12 tablets in 24 hours, which is the maximum daily amount • child takes more than 5 doses in 24 hours • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product	
	<b>Do not use</b> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	

**HSINIVA NO VARNISH**



<b>Drug Facts (continued)</b>
• If you are allergic to acetaminophen or any of the inactive ingredients in this product.
<b>Ask a doctor before use if you have liver disease.</b>
<b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</b>
<b>Stop use and ask a doctor if</b>
• Pain gets worse or lasts more than 10 days in adults and children.
• Pain gets worse or lasts more than 5 days in children under 12 years.
• Fever gets worse or lasts more than 3 days.
• New symptoms occur.
• Redness or swelling is present.
These could be signs of a serious condition.
<b>If pregnant or breast-feeding</b> , ask a health professional before use.
<b>Keep out of reach of children.</b>
<b>Overdose warning:</b> Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
<b>Directions</b>
<b>Do not take more than directed</b>
<b>Adults and Children 12 years and over.</b>
• Take 2 tablets every 4 to 6 hours while symptoms last.
• Do not take more than 12 tablets in 24 hours
<b>Children 6-11 years:</b>
• Take 1 tablet every 4 to 6 hours while symptoms last.
• Do not take more than 5 tablets in 24 hours
<b>Children under 6 years:</b>
Do not use this adult regular strength product in children under 6 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and may cause liver damage.
<b>Other information</b>
• Store between 20-25°C (68-77°F).
<b>Inactive ingredients</b> Pregelatinized Starch, Polydome K-30, Sodium starch Glycolate, Stearic acid.
<b>STOP PEELING</b>

**ACETAMINOPHEN 325MG**

acetaminophen tablet

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:10135-828
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg	
Inactive Ingredients				
Ingredient Name			Strength	
POVIDONE K30 (UNII: U725QWY32X)				
STARCH, CORN (UNII: O8232NY3SJ)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	C;19	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10135-828-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2026	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	02/01/2026		

**Labeler** - Marlex Pharmaceuticals Inc (782540215)

Revised: 2/2026

Marlex Pharmaceuticals Inc