

**REINREUDE PROVIDES TEMPORARY RELIEF FOR MINOR MUSCLE AND JOINT PAIN- lidocaine 4% provides temporary relief for minor muscle and joint pain patch**  
**Jiangxi Yudexi Pharmaceutical Co., LTD**

-----  
**85248-076**

**Active Ingredient**

Lidocaine 4%

**Purpose**

Pain Relief

**Use**

temporarily relieves minor pain associated with: arthritis simple backache bursitis tendonitis muscle strains sprains bruises cramps

**Warnings**

For external use only

**Do not use**

use only as directed. Read and follow all directions and warnings on this carton. avoid contact with eyes and mucous membranes rare cases of serious burns have been reported with products of this type do not apply to wounds or damaged skin, broken or irritated skin do not bandage tightly or apply local heat (such as heating pads) to the area of use A transient burning sensation may occur upon application but usually disappears within a few days.

**When Using**

use only as directed. Read and follow all directions and warnings on this carton. avoid contact with eyes and mucous membranes rare cases of serious burns have been reported with products of this type do not apply to wounds or damaged skin, broken or irritated skin do not bandage tightly or apply local heat (such as heating pads) to the area of use A transient burning sensation may occur upon application but usually disappears within a few days.

**Stop Use**

condition worsens redness is present irritation develops symptoms persist for more than 7 days or clear up and occur again within a few dsys you experience signs

of skin injury, such as pain, swelling, or blistering where the product was applied. If pregnant or breast-feeding, ask a health professional before use.

### **Ask Doctor**

■ condition worsens ■ redness is present ■ irritation develops ■ symptoms persist for more than 7 days or clear up and occur again within a few days ■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

### **Keep Out Of Reach Of Children**

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

adults and children over 12 years. ■ remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle ■ carefully remove smaller portion of backing from patch and apply exposed portion of patch to affected area ■ once exposed portion of patch is positioned, carefully remove remaining backing to completely apply patch to affected area ■ Wear one patch for up to 8 hours. ■ Repeat if needed, but do not use more than 3 times daily. children 12 years or younger: ask a doctor

### **Inactive ingredients**

water, aluminum hydroxide, cellulose gum, glycerin, isopropyl myristate, nonoxonyl-30, polyacrylic acid, polysorbate 80, sodium polyacrylate, sorbitan sesquioleate, Starch, talc, tartaric acid, titanium dioxide

### **PRINCIPAL DISPLAY PANEL**

B01.621

TEMPORARILY RELIEVES  
MINOR ACHES & PAINS


  

  

**TEMPORARILY RELIEVES**  
**MINOR ACHES & PAINS**

**BACK, NECK,  
LEG & SHOULDER**

PROVIDES TEMPORARY RELIEF  
FOR MINOR MUSCLE AND JOINT  
PAIN  
TOPICAL ANESTHETIC

- No-Mess, Single-Use Application
- Easy to Apply and Remove
- Flexible Gel Patch Designed For Secure Application
- Unscented

**8 PATCHES**



BACK, NECK, LEG & SHOULDER

**Drug Facts**

Active Ingredient..... Purpose  
Lidocaine 4%..... Pain Relief

**Uses**  
temporarily relieves minor pain associated with:  
arthritis simple backache bursitis tendonitis muscle strains sprains bruises cramps

**Warnings**  
For external use only

**When using this product**  
use only as directed. Read and follow all directions and warnings on this carton.  
avoid contact with eyes and mucous membranes  
rare cases of serious burns have been reported with products of this type  
do not apply to wounds or damaged skin, broken or irritated skin  
do not bandage tightly or apply local heat (such as heating pads) to the area of use  
A transient burning sensation may occur upon application but usually disappears within a few days.

**Stop use and ask a doctor if** condition worsens redness is present irritation develops  
symptoms persist for more than 7 days or clear up and occur again within a few days  
you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

**If pregnant or breast-feeding** ask a health professional before use.  
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
adults and children over 12 years.  
remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle  
carefully remove smaller portion of backing from patch and apply exposed portion of patch to affected area  
once exposed portion of patch is positioned, carefully remove remaining backing to completely apply patch to affected area  
Wear one patch for up to 8 hours.  
Repeat if needed, but do not use more than 3 times daily.  
children 12 years or younger: ask a doctor

**Inactive Ingredients**  
water, aluminum hydroxide, cellulose gum, glycerin, isopropyl myristate, nonoxonyl-30, polyacrylic acid, polysorbate 80, sodium polyacrylate, sorbitan sesquioleate, Starch, talc, tartaric acid, titanium dioxide.




DISTRIBUTOR: JIANGXI YUDEKI PHARMACEUTICAL CO., LTD  
ADDRESS: NO.1998, JINGKAI WEST 5TH ROAD, CHENGBEI  
JINGKAI DISTRICT, ZHANGSHU CITY, YICHUN CITY,  
JIANGXI PROVINCE



6 979843 980678

## REINREUDE PROVIDES TEMPORARY RELIEF FOR MINOR MUSCLE AND JOINT PAIN

lidocaine 4% provides temporary relief for minor muscle and joint pain patch

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:85248-076
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100

### Inactive Ingredients

Ingredient Name	Strength
SORBITAN SESQUIOLEATE (UNII: 0W8RRI5W5A)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>NONOXYNOL-30</b> (UNII: JJX07DG188)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)
<b>TARTARIC ACID</b> (UNII: W4888I119H)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>CELLULOSE GUM</b> (UNII: K679OBS311)
<b>SODIUM POLYACRYLATE (250000 MW)</b> (UNII: 05I15JNI2J)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>POLYACRYLIC ACID (250000 MW)</b> (UNII: 9G2MAD7J6W)

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85248-076-01	8 in 1 BOX; Type 0: Not a Combination Product	02/08/2026	

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/08/2026	

**Labeler** - Jiangxi Yudexi Pharmaceutical Co., LTD (455662836)

Revised: 2/2026

Jiangxi Yudexi Pharmaceutical Co., LTD