

LIDOCAINE PAIN RELIEF GEL-PATCH- lidocaine 4% patch
Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.

LIDOCAINE
PAIN RELIEF GEL-PATCH

Active ingredient

Lidocaine 4% Purpose: Topical Anesthetic

Purpose

Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Ask a doctor if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

Stop use if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children

Keep out of reach of children. If swallowed accidentally , get medical help or contact with Poison Control Center immediately.

Directions**Adults and Children above 12 years old:**

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under age of 12: Consult a doctor

Other information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, PropyleneGlycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments?

supports@povihomeofficial.com

Package label. Principal display panel

Povihome®
The more attention to health

Shoulder
Knee
Back

LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomeofficial.com

Maximum OTC Strength
LIDOCAINE
PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

Odor-Free

Single-Use Application

Numbs Pain Away

15 (PATCHES)
5.51" X 3.93"
14cm X 10cm

L PATCH
Lidocaine 4%

Usage

- Remove patch from outer package
- Partially peel back protective film
- Place on affected area
- Remove remaining film while pressing patch to skin

Drug Facts

Active ingredient	Purpose
Lidocaine 4%	Topical anesthetic

Warnings
For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or malleae
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- severe skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,
ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.
If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions
Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under age of 12: Consult a doctor

Other Information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive Ingredients
Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, PropyleneGlycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:
supports@povihomeofficial.com

Child Resistant Packaging

6 0 2 4 2 6 8 0 1 5 5 2

Guangzhou Luquanmao Import & Export Trade Co., Ltd.
412, Building A, No. 6, Dong Heng San Road,
Zhucun, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC



LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomemedical.com

Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

Odor-Free

Single-Use Application

Numbs Pain Away



16 PATCHES
5.51" X 3.93"
14cm X 10cm

L PATCH
Lidocaine 4%

Guangzhou Liaqunmao Import & Export Trade Co., Ltd.
417, Building A, No. 8, Dong Heng San Road,
Zhuosen, Taishan District, Guangzhou, Guangdong, China

Made in PRC

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

The more attention to health
Povihome®

Drug Facts

Active ingredient Purpose
Lidocaine 4% Topical anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under age of 12: Consult a doctor

Other information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polyacrylate G9, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:

support@povihomemedical.com

Child Resistant Packaging





Shoulder



Knee



Back

LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomeofficial.com

Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

Odor-Free

Single-Use Application

Numbs Pain Away



29 PATCHES
5.51" X 3.93"
14cm X 10cm

L PATCH
Lidocaine 4%

Guangzhou Lidoqunman Import & Export Trade Co., Ltd.
412 Building A, No. 6, Dongyuan Street, San Ping,
Zhuosen, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

Drug Facts

Active ingredient Purpose
Lidocaine 4% Topical anesthetic

Uses
Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hip, shoulder, elbow.

Warnings
For external use only

Do not use
• more than 1 patch on your body at a time or with other topical anesthetics of the same time
• on wounds or damaged skin
• with a heating pad or device
• with other ointments, creams, sprays, or liniments
• if you are allergic to any ingredients of this product

When using this product
• use only as directed
• avoid contact with the eyes, mucous membranes, or mouth
• do not bandage tightly
• dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests the patch.

Stop use and ask a doctor if
• localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
• condition worsens
• symptoms persist for more than 7 days
• symptoms clear up and occur again within a few days

The more attention to health
Povihome®

If pregnant or breast-feeding,
ask a health professional before use.
KEEP OUT OF REACH OF CHILDREN.
If swallowed accidentally, get medical help or contact with Poison Control Center immediately

Directions
Adults and Children above 12 years old:
• Clean and dry the affected area.
• Tear off the protective film and apply the exposed part of the patch to the affected area.
• Carefully remove remaining film while pressing the patch firmly on the skin.
• Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
• Remove patch from the skin after at most 8 hour application
Children under age of 12: Consult a doctor

Other information
• Store at 20°C to 25°C (68°F - 77°F)
• Store in a cool dry place away from direct sunlight

Inactive ingredients
Dihydroxyacetone, Ammonium Chloride, Ethanol, Disodium Citrate, Glycerin, Kaolin, Methylparaben, Polyacrylate Gels, Propylene Glycol, Polysorbate Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Terephthalic Acid, Water

Questions or Comments:
support@povihomeofficial.com

Child Resistant Packaging





LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomemedical.com

Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

- Odor-Free
- Single-Use Application
- Numbs Pain Away



31 (PATCHES)
5.51" X 3.93"
14cm X 10cm

L PATCH
Lidocaine 4%

Shanghai Povihome Import & Export Trade Co., Ltd.
411 Building A, No. 8, Dong Heng San Road,
Zhongchuan, Xuhui District, Shanghai, Shanghai, China

Model: L-P101

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

Drug Facts

Active ingredient Purpose
Lidocaine 4% Topical anesthetic

Uses

Temporarily relieves muscle aches and minor joint pain in the wrist, knee, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or lotions
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or mouth
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests the patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as red, itching, redness, itching, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application.

Children under age of 12: Consult a doctor

Other information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients

Dipropylamine, Dimethylacetamide, Dibutyl Tin Diolate, Glycerin, Acetic Acid, Methylparaben, Polyacrylate G1, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:

Support@povihomemedical.com

Child Resistant Packaging



6 9324 10010 1 001

Povihome®
The more attention to health

LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomeofficial.com



Povihome®
The more attention to health.

Maximum OTC Strength

LIDOCAINE
PAIN RELIEF GEL-PATCH

Desensitizes Aggravated
Nerves And Relieves Pain.

- Odor-Free
- Single-Use Application
- Numbs Pain Away



35 PATCHES
5.51" X 3.93"
14cm X 10cm

L PATCH
Lidocaine 4%

Shanghai Liuzhenmao Import & Export Trade Co., Ltd.
111 Building, No. 6, Dong Feng San Road,
Shanghai, China

Usage



Remove patch from
outer package



Partially peel back
protective film



Place on
affected area




Remove remaining film
while pressing patch to skin

The more attention to health.
Povihome®

Drug Facts		If pregnant or breast-feeding, ask a health professional before use.
Active ingredient Purpose Lidocaine 4% Topical anesthetic.		KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.
Uses Temporarily relieves muscle aches and minor joint pains in the wrist, knee, back, neck, hips, shoulders, elbows.		Directions Adults and Children above 12 years old: • Clean and dry the affected area. • Tear off the protective film and apply the exposed part of the patch to the affected area. • Carefully remove remaining film while pressing the patch firmly on the skin. • Apply 1 patch at a time to affected area, not more than 3 to 4 times daily. • Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.
Warnings For external use only Do not use • more than 1 patch on your body at a time or with other topical anesthetics at the same time. • on wounds or damaged skin. • with a heating pad or device. • with other ointments, creams, sprays, or liniments. • if you are allergic to any ingredients of this product. When using this product • use only as directed. • avoid contact with the eyes, mucous membranes, or mucosa. • do not bandage tightly. • dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests the patch. Stop use and ask a doctor if • localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling, and blistering condition worsens. • symptoms persist for more than 7 days. • symptoms clear up and occur again within a few days.	Other Information • Store at 20°C to 25°C (68°F - 77°F). • Store in a cool dry place away from direct sunlight. Inactive Ingredients Chlorophthalmum, Anticollinesterase, Estrone Diolium, Quercetin, Kavaol, Methylparaben, Polysorbate 80, Propylparaben, Polypropylene Glycol, Polyvinylpyrrolidone K90, Propylparaben, Sodium Phosphate, Titanium Dioxide, Talcum, ACAC, water.	
Questions or Comments: support@povihomeofficial.com		Child Resistant Packaging





LIDOCAINE PAIN RELIEF GEL-PATCH

lidocaine 4% patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83602-257
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.04 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

GLYCERIN (UNII: PDC6A3C0OX)
POVIDONE K90 (UNII: RDH86HJV5Z)
METHYLPARABEN (UNII: A2I8C7HI9T)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
EDETATE DISODIUM (UNII: 7FLD91C86K)
KAOLIN (UNII: 24H4NWX5CO)
TARTARIC ACID (UNII: W4888I119H)
WATER (UNII: 059QF0K0OR)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83602-257-15	15 in 1 BOX	02/09/2026	
1		1 in 1 BAG		
1	NDC:83602-257-01	12 g in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:83602-257-16	16 in 1 BOX	02/09/2026	
2		1 in 1 BAG		
2	NDC:83602-257-01	12 g in 1 PATCH; Type 0: Not a Combination Product		
3	NDC:83602-257-31	31 in 1 BOX	02/09/2026	
3		1 in 1 BAG		
3	NDC:83602-257-01	12 g in 1 PATCH; Type 0: Not a Combination Product		
4	NDC:83602-257-35	35 in 1 BOX	02/09/2026	
4		1 in 1 BAG		
4	NDC:83602-257-01	12 g in 1 PATCH; Type 0: Not a Combination Product		
5	NDC:83602-257-36	36 in 1 BOX	02/09/2026	
5		1 in 1 BAG		
5	NDC:83602-257-01	12 g in 1 PATCH; Type 0: Not a Combination Product		
6	NDC:83602-257-29	29 in 1 BOX	02/09/2026	
6		1 in 1 BAG		

6 NDC:83602-257-01 12 g in 1 PATCH; Type 0: Not a Combination Product

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/09/2026	

Labeler - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Registrant - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Revised: 2/2026

Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.