

MEIJER - avobenzone, homosalate, octisalate, octocrylene, oxybenzone lotion
Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredients

Avobenzone 3.0%
Homosalate 13.0%
Octisalate 5.0%
Octocrylene 7.0%
Oxybenzone 4.0%

Purpose

Sunscreen

Uses

- Helps prevent sunburn
- If used as directed with other sun protection measures (see **Directions**), decrease the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only. Do not use on damaged or broken skin. May stain some fabrics.

When using this product

Keep out of eyes. Rinse with water to remove.

stop use and ask a doctor

if rash occurs

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply liberally 15 minutes before sun exposure
- reapply
- after 80 minutes of swimming or sweating
- immediately after towel drying
- at least every 2 hours
- **Sun Protection Measures:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
- limit time in the sun, especially from 10 a.m. - 2 p.m.

- wear long-sleeve shirts, pants, hats and sunglasses
- children under 6 months of age: Ask a doctor

Inactive Ingredients


Water, Sorbitol, Stearic Acid, Triethanolamine, Aluminum Starch Octenylsuccinate, Benzyl Alcohol, Sorbitan Isostearate, VP/Eicosene Copolymer, Dimethicone, Polyglyceryl-3 Distearate, Carbomer, Tocopherol, Disodium EDTA, Methylparaben, Propylparaben, Fragrance.

Other Information

Protect this product from excessive heat and direct sun.

Principle Display Panel



Drug Facts	
Active Ingredients	Purpose
Avobenzone 3.0%, Homosalate 13.0%, Octisalate 5.0%, Octocrylene 7.0%, Oxybenzone 4.0%	Sunscreen
Uses • helps prevent sunburn • if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun	
Warnings For external use only • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask a doctor if rash occurs • Keep out of the reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. • May stain some fabrics.	
Directions • apply liberally 15 minutes before sun exposure • reapply: after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m. • wear long-sleeve shirts, pants, hats and sunglasses • children under 6 months of age: Ask a doctor	
Inactive Ingredients Water, Sorbitol, Stearic Acid, Triethanolamine, Aluminum Starch Octenylsuccinate, Benzyl Alcohol, Sorbitan Isostearate, VP/Eicosene Copolymer, Dimethicone, Polyglyceryl-3 Distearate, Carbomer, Tocopherol, Disodium EDTA, Methylparaben, Propylparaben, Fragrance.	
Other information • protect this product from excessive heat and direct sun Made in USA	
	
DIST. BY MEIJER DISTRIBUTION, INC. 2929 WALKER AVE NW GRAND RAPIDS, MI 49544 www.meijer.com	
90005553	

Principal Display Panel

.25" NO PR

meijer

Sunscreen lotion

hypoallergenic
UVA/UVB Protection

50

Broad Spectrum
SPF 50

Water Resistant
(80 Minutes)

3 FL. OZ. (88 mL)

.25" NO PR

Drug Facts

Active ingredients Avobenzone 3.0%, Homosalate 13.0%, Octisalate 5.0%, Octocrylene 7.0%, Oxybenzone 4.0%	Purpose Sunscreen
--	-----------------------------

Uses • helps prevent sunburn • if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings
For external use only • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask a doctor if rash occurs • Keep out of the reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. • May stain some fabrics

Directions
• apply liberally 15 minutes before sun exposure
• reapply: after 80 minutes of swimming or sweating
• immediately after towel drying • at least every 2 hours
• **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m. • wear long-sleeve shirts, pants, hats and sunglasses
• children under 6 months of age: Ask a doctor

Inactive ingredients Water, Sorbitol, Stearic Acid, Triethanolamine, Aluminum Starch Octenylsuccinate, Benzyl Alcohol, Sorbitan Isostearate, VP/Eicosene Copolymer, Dimethicone, Polyglyceryl-3 Distearate, Carbomer, Tocopherol, Disodium EDTA, Methylparaben, Propylparaben, Fragrance.

Other information • protect this product from excessive heat and direct sun



Made in USA



DIST BY MEIJER
DISTRIBUTION, INC.
2929 WALKER AVE NW
GRAND RAPIDS, MI 49544
www.meijer.com 93005374

Principal Display Panel



Drug Facts	
Active ingredients	Purpose
Avobenzone 3.0%, Homosalate 13.0%, Octisalate 5.0%, Octocrylene 7.0%, Oxibenzonone 4.0%	Sunscreen
Uses • helps prevent sunburn • if used as directed with other sun protection measures (see <i>Directions</i>), decreases the risk of skin cancer and early skin aging caused by the sun	
Warnings For external use only • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask a doctor if rash occurs • Keep out of the reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. • May stain some fabrics.	
Directions • apply liberally 15 minutes before sun exposure • reapply: after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m. • wear long-sleeve shirts, pants, hats and sunglasses • children under 6 months of age: Ask a doctor	
Inactive ingredients Water, Sorbitol, Stearic Acid, Triethanolamine, Aluminum Starch Octenylsuccinate, Benzyl Alcohol, Sorbitan Isostearate, VP/Eicosene Copolymer, Dimethicone, Polyglyceryl-3 Distearate, Carbomer, Tocopherol, Disodium EDTA, Methylparaben, Propylparaben, Fragrance.	
Other information • protect this product from excessive heat and direct sun	
 <p>Made in USA DST. BY MEIJER DISTRIBUTION, INC. 7329 WALKER AVE NW GRAND RAPIDS, MI 49544 www.meijer.com</p> 	

MEIJER

avobenzone, homosalate, octisalate, octocrylene, oxybenzone lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-006
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 mL in 100 mL
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	13 mL in 100 mL

OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	5 mL in 100 mL
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	7 mL in 100 mL
OXYBENZONE (UNII: 950OS7VE0Y) (OXYBENZONE - UNII:950OS7VE0Y)	OXYBENZONE	4 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TROLAMINE (UNII: 9O3K93S3TK)	
ALUMINUM STARCH OCTENYLSUCCINATE (UNII: I9PJ0O6294)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
SORBITAN ISOSTEARATE (UNII: 01S2G2C1E4)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
TOCOPHEROL (UNII: R0ZB2556P8)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-006-57	295 mL in 1 BOTTLE		
2	NDC:41250-006-22	88 mL in 1 TUBE		
3	NDC:41250-006-19	473 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	01/01/2012	

Labeler - Meijer Distribution Inc (006959555)