

**FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE -
fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film
coated, extended release
AmeriSource Bergen**

Fexofenadine Hydrochloride and Pseudoephedrine Hydrochloride

Drug Facts

<i>Active ingredients (in each extended-release tablet)</i>	<i>Purpose</i>
Fexofenadine HCl, USP 60 mg	Antihistamine
Pseudoephedrine HCl, USP 120 mg	Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland

- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- **do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1800-222-1222).

Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- do not use if carton is opened or if individual blister units are torn or opened
- store between 20° to 25°C (68° to 77°F)
- Meets dissolution test 6

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, ethyl cellulose, ferric oxide yellow, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid.

Questions or comments?

Call toll-free **1-800-818-4555 weekdays**

Distributed By:
AmerisourceBergen
1 West First Avenue
Conshohocken, PA 19428

PRINCIPAL DISPLAY PANEL - 30 Tablet Blister Pack Carton

Compare to Allegra-D[®] active ingredients**

GOOD
NEIGHBOR
PHARMACY[®]

NDC 46122-694-65

NON-DROWSY

12 HOUR

fexofenadine HCl 60 mg/antihistamine
pseudoephedrine HCl 120 mg/nasal decongestant
Extended-Release Tablets, USP


Indoor & Outdoor Allergies

Allergy & Congestion

Indoor/Outdoor Relief of:

- Nasal and Sinus Congestion Due to Colds or Allergies
- Sneezing; Runny Nose; Itchy, Watery Eyes and Itchy Nose or Throat Due to Allergies

30 Extended-Release Tablets



Drug Facts (continued)

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1800-222-1222).

Drug Facts (continued)

Directions

do not divide, crush, chew or dissolve the tablet; swallow tablet whole	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
adults and children 12 years of age and over	do not use
children under 12 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- do not use if carton is opened or if individual blister units are torn or opened
- store between 20° to 25°C (68° to 77°F)
- Meets dissolution test 6

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, ethyl cellulose, ferric oxide yellow, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid.

Questions or comments?
Call toll-free 1-800-818-4355 weekdays

5265344

DO NOT USE IF INDIVIDUAL BLISTER UNIT IS OPEN OR TORN

REV-08/20

3 46122 69455 0

AB# 10293617

DNDH/PHUS/NH/138

MADE IN INDIA

Quibora or © 2020 N7VZ www.n7vp.com

CONTRACT MAN, PA 19428
1 West First Avenue
Aurora, IL 60009

Dist. by:

fexofenadine HCl 60 mg/antihistamine
pseudoephedrine HCl 120 mg/nasal decongestant
Extended-Release Tablets, USP

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Allegra-®.

30 Extended-Release Tablets

fexofenadine HCl 60 mg/antihistamine
pseudoephedrine HCl 120 mg/nasal decongestant
Extended-Release Tablets, USP

Allergy & Congestion

GOOD NEIGHBOR PHARMACY

Compare to Allegra-® active ingredients

NDC 46122-694-06

NON-DROWSY 12 HOUR

fexofenadine HCl 60 mg/antihistamine
pseudoephedrine HCl 120 mg/nasal decongestant
Extended-Release Tablets, USP

Allergy & Congestion

Indoor/Outdoor Relief of:

- Nasal and Sinus Congestion Due to Colds or Allergies
- Sneezing; Runny Nose; Itchy, Watery Eyes and Itchy Nose or Throat Due to Allergies

30 Extended-Release Tablets

**NON VARNISH
NO INK AREA**

6 x 5 Tablets

Drug Facts

Active ingredients (in each extended-release tablet)	Purpose
Fexofenadine HCl, USP 60 mg.....	Antihistamine
Pseudoephedrine HCl, USP 120 mg.....	Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages

Drug Facts (continued)

- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE

fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-694
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FEXOFENADINE HYDROCHLORIDE (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	60 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
POVIDONE K30 (UNII: U725QWY32X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	white, yellow	Score	no score
Shape	CAPSULE (bilayer)	Size	17mm
Flavor		Imprint Code	724
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-694-65	3 in 1 CARTON	03/01/2018	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090818	03/01/2018	

Labeler - AmeriSource Bergen (007914906)

Establishment

Name	Address	ID/FEI	Business Operations
Sun Pharmaceutical Industries Limited		650445203	analysis(46122-694) , manufacture(46122-694)

Revised: 11/2025

AmeriSource Bergen