

**BASIC CARE DAYTIME SEVERE COLD AND FLU NIGHTTIME SEVERE COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl
Amazon.com Services LLC**

Amazon Daytime Severe Cold & Flu Nighttime Severe Cold & Flu Drug Facts

**Nighttime Severe Cold & Flu
Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children

- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Daytime Severe Cold & Flu

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In care of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

amazon

basic care

Multi-Symptom Relief

Compare to Vicks® DayQuil® Severe active ingredients

Non-Drowsy

Daytime Severe Cold & Flu

Acetaminophen

Dextromethorphan HBr

Guaifenesin

Phenylephrine HCl

Pain Reliever/Fever Reducer

Nasal Decongestant

Cough Suppressant

Expectorant

ALCOHOL FREE

Max Strength

Original Flavor

Maximum Strength

Compare to Vicks[®] NyQuil[®] Severe active ingredients

Nighttime Severe Cold & Flu

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

Pain Reliever, Fever Reducer, Nasal Decongestant, Cough Suppressant, Antihistamine

ALCOHOL FREE

Berry Flavor

COMBINATION PACK

2 - 12 FL OZ (355 mL) BOTTLES, TOTAL 24 FL OZ (1.5 pt) 710 mL

amazon

basic+care

Daytime
Severe
Cold & Flu

Nighttime
Severe
Cold & Flu

amazon

basic+care

Multi-Symptom
Relief

Compare to
Vicks® DayQuil®
Severe
active ingredients*

Maximum
Strength

NDC 72288-597-02

Compare to
Vicks® NyQuil®
Severe
active ingredients*

Non-Drowsy

Daytime
Severe
Cold & Flu

Acetaminophen
Dextromethorphan HBr
Guaifenesin
Phenylephrine HCl

Pain Reliever/Fever Reducer
Nasal Decongestant
Cough Suppressant
Expectorant

ALCOHOL FREE
Max Strength

Original
Flavor

Nighttime
Severe
Cold & Flu

Acetaminophen,
Phenylephrine HCl,
Doxylamine Succinate,
Dextromethorphan HBr

Pain Reliever, Fever Reducer,
Nasal Decongestant,
Cough Suppressant,
Antihistamine

ALCOHOL FREE



Berry
Flavor

COMBINATION PACK

2 - 12 FL OZ (355 mL) BOTTLES,
TOTAL 24 FL OZ (1.5 pt) 710 mL

Nighttime Relief

Headache, Fever,
Sore Throat,
Minor Aches & Pains,
Nasal/Sinus Congestion
& Sinus Pressure,
Sneezing, Runny Nose,
Cough



DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING

DISTRIBUTED BY:
AMAZON.COM SERVICES LLC
410 TERRY AVENUE N.
SEATTLE, WA 98109


© 2025 AMAZON.COM, INC. OR
ITS AFFILIATES. ALL RIGHTS
RESERVED. BASIC CARE AND
ALL RELATED LOGOS ARE
TRADEMARKS OF AMAZON.COM,
INC. OR ITS AFFILIATES.



PAPER | PLASTIC | PLASTIC | 2
BOX | BOTTLE | CUP | 01/17

3 70030 14549 2

59702 BN C3



Nighttime Severe Cold & Flu

Daytime Severe Cold & Flu

Drug Facts	
Active ingredients (in each 30 mL)	Purpose
Acetaminophen 660 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

Drug Facts	
Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if:
 ■ adult takes more than 4,000 mg of acetaminophen in 24 hours
 ■ child takes more than 5 doses in 24 hours
 ■ taken with other drugs containing acetaminophen
 ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have
 ■ liver disease ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes ■ glaucoma
 ■ cough that occurs with too much phlegm (mucus)
 ■ a breathing problem such as an emphysema or chronic bronchitis
 ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
 ■ a sodium-restricted diet

Ask a doctor before use if you have
 ■ liver disease ■ heart disease ■ thyroid disease
 ■ diabetes ■ high blood pressure
 ■ trouble urinating due to an enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus)
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are
 ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
 When using this product do not use more than directed

When using this product
 ■ do not use more than directed ■ excitability may occur, especially in children
 ■ marked drowsiness may occur ■ avoid alcoholic drinks
 ■ be careful when driving a motor vehicle or operating machinery
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
 ■ you get nervous, dizzy or sleepless
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days (children) or 7 days (adults)
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts.
 These could be signs of a serious condition.

Stop use and ask a doctor if
 ■ you get nervous, dizzy or sleepless
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts.
 These could be signs of a serious condition.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed – see Overdose warning
 ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Directions

■ take only as directed – see Overdose warning
 ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information
 ■ each 30 mL contains: sodium 41 mg
 ■ store at 20-25°C (68-77°F)

Other information
 ■ each 15 mL contains: sodium 6 mg
 ■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9260

Questions or comments? 1-800-719-9260

Daytime Relief

Headache, Fever, Sore Throat, Minor Aches & Pains, Nasal/Sinus Congestion & Sinus Pressure, Chest Congestion, Cough



*These products are not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Severe and Vicks® NyQuil® Severe.

GLUTEN FREE

PARENTS:
 Learn about teen medicine abuse

www.StopMedicineAbuse.org



BASIC CARE DAYTIME SEVERE COLD AND FLU NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72288-597
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-597-02	1 in 1 CARTON; Type 0: Not a Combination Product	03/11/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

Part 1 of 2

BASIC CARE NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:72288-639
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	RED (clear)	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-639-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Part 2 of 2

BASIC CARE DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:72288-603
---------------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	ORANGE (clear)	Score	
Shape		Size	
Flavor	FRUIT, MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-603-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/11/2021	

Labeler - Amazon.com Services LLC (128990418)

Revised: 10/2025

Amazon.com Services LLC