

**REVASOL ACETAMINOPHEN SUSPENSION DYE FREE- acetaminophen liquid
Rnv LLC**

Revasol Acetaminophen Suspension Grape Flavor Dye Free. NDC 84379-357-16

Acetaminophen 160 mg

Pain reliever/fever reducer

Uses

• temporarily: • reduces fever • relieves minor aches and pains due to: • the common cold • headache • toothache • backache • premenstrual and menstrual cramps • flu • sore throat • muscular aches • minor pain of arthritis

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

• adult

takes more than 4,000 mg of acetaminophen in 24 hours • child takes more than 5 doses in 24

hours, which is the maximum daily amount • taken with other drugs containing acetaminophen

• adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or

followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not

sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to

acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days in adults

• pain

gets worse or lasts more than 5 days in children under 12 years • fever gets worse or lasts

more than 3 days • new symptoms occur • redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs of symptoms.

Directions

• **do not give more than directed (see overdose warning)** • shake well before using • mL=milliliter • **Children 2 years to under 12 years:** • find right dose on chart below: If possible use weight to dose, otherwise, use age • remove the child protective cap and squeeze the dose into the dosing cup • repeat dose every 4 hours while symptoms last • do not take more than 5 times in 24 hours unless directed by a doctor • do not take for more than 5 days unless directed by a doctor

Weight (lb)	Age (years)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-85 lbs	11 years	15 mL

*or as directed by a doctor

• **Adults and children 12 years and over:** • take 20 mL (2 x 10 mL) every 4 hours while symptoms last. For adult dosing, the dosing cup needs to be filled to 10 mL line twice. • do not take more than 5 doses (100 mL) in 24 hours unless directed by a doctor • do not take for more than 10 days unless directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product . Do not use any other dosing device.

Other information

- store between 15-30°C (59-86°F).
- Do not use if printed safety seal is torn, broken or missing.

Inactive ingredients Anhydrous citric acid, carboxymethylcellulose, flavors, glycerin, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

Questions or comments? 1-888-446-4753

Monday - Friday (9 AM - 5 PM EST)

NDC 84379-357-16



ACETAMINOPHEN SUSPENSION

Acetaminophen 160 mg in 5 mL

Oral Suspension Pain Reliever and
Fever Reducer Age 2 years to adults

Grape Flavor
Dye Free

16 FL OZ (474 mL)

Drug Facts																						
Active Ingredient (in each 5 mL)	Purpose																					
Acetaminophen 160 mg	Pain reliever/fever reducer																					
Uses • temporarily • reduces fever • relieves minor aches and pains due to: • the common cold • headache • toothache • backache • premenstrual and menstrual cramps • flu • sore throat • muscular aches • minor pain of arthritis																						
Warnings																						
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if • adult takes more than 4,000 mg of acetaminophen in 24 hours • child takes more than 5 doses in 24 hours, which is the maximum daily amount • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions.</p> <p>Symptoms may include: • skin reddening • blisters • rash</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to acetaminophen or any of the inactive ingredients in this product <p>Ask a doctor before use if you have liver disease</p> <p>Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin</p> <p>When using this product do not exceed recommended dose (see overdose warning)</p> <p>Stop use and ask a doctor if • pain gets worse or lasts more than 10 days in adults • pain gets worse or lasts more than 5 days in children under 12 years • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present</p> <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of the reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs of symptoms.</p>																						
Directions • do not give more than directed (see overdose warning) • shake well before using • mL=milliliter • Children 2 years to under 12 years: • find right dose on chart below. If possible use weight to dose, otherwise, use age • remove the child protective cap and squeeze the dose into the dosing cup • repeat dose every 4 hours while symptoms last • do not take more than 5 times in 24 hours unless directed by a doctor • do not take for more than 5 days unless directed by a doctor																						
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Weight (lb)</th> <th>Age (years)</th> <th>Dose (mL)*</th> </tr> </thead> <tbody> <tr> <td>under 24</td> <td>under 2 years</td> <td>ask a doctor</td> </tr> <tr> <td>24-35 lbs</td> <td>2-3 years</td> <td>5 mL</td> </tr> <tr> <td>36-47 lbs</td> <td>4-5 years</td> <td>7.5 mL</td> </tr> <tr> <td>48-59 lbs</td> <td>6-8 years</td> <td>10 mL</td> </tr> <tr> <td>60-71 lbs</td> <td>9-10 years</td> <td>12.5 mL</td> </tr> <tr> <td>72-85 lbs</td> <td>11 years</td> <td>15 mL</td> </tr> </tbody> </table> <p>*or as directed by a doctor</p> <p>• Adults and children 12 years and over: take 20 mL (2 x 10 mL) every 4 hours while symptoms last. For adult dosing, the dosing cup needs to be filled to 10 mL line twice. • do not take more than 5 doses (100 mL) in 24 hours unless directed by a doctor • do not take for more than 10 days unless directed by a doctor</p> <p>Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.</p>		Weight (lb)	Age (years)	Dose (mL)*	under 24	under 2 years	ask a doctor	24-35 lbs	2-3 years	5 mL	36-47 lbs	4-5 years	7.5 mL	48-59 lbs	6-8 years	10 mL	60-71 lbs	9-10 years	12.5 mL	72-85 lbs	11 years	15 mL
Weight (lb)	Age (years)	Dose (mL)*																				
under 24	under 2 years	ask a doctor																				
24-35 lbs	2-3 years	5 mL																				
36-47 lbs	4-5 years	7.5 mL																				
48-59 lbs	6-8 years	10 mL																				
60-71 lbs	9-10 years	12.5 mL																				
72-85 lbs	11 years	15 mL																				
Other information																						
<ul style="list-style-type: none"> • store between 15-30°C (59-86°F). • Do not use if printed safety seal is torn, broken or missing. <p>Inactive ingredients Anhydrous citric acid, carboxymethylcellulose, flavors, glycerin, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.</p>																						
Questions or comments? 1-888-446-4753 Monday - Friday (9 AM - 5 PM EST)																						
<p>Made in USA Distributed by: RNV LLC 14501 NW 60 Ave, Miami Lakes, FL 33014</p>																						



8 50073 25715 9

Lot: _____
Rev. 1 Exp: _____

REVASOL ACETAMINOPHEN SUSPENSION DYE FREE

acetaminophen liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84379-357
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
XANTHAN GUM (UNII: TTV12P4NEE)	
WATER (UNII: 059QF0KO0R)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	Score
Shape	Size

Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84379-357-16	474 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/01/2025	

Labeler - Rnv LLC (118917568)

Registrant - Rnv LLC (118917568)

Establishment

Name	Address	ID/FEI	Business Operations
Rnv LLC		118917568	manufacture(84379-357)

Revised: 2/2026

Rnv LLC