

**ADVANTAGE MULTI FOR CATS- imidacloprid, moxidectin solution**  
**Elanco US Inc.**

-----  
**Elanco™**  
**advantage multi™**  
**for cats**  
**(imidacloprid+moxidectin)**  
**Topical Solution**

**Once-a-month topical solution for cats for the prevention of heartworm disease, kills adult fleas, is indicated for the treatment of flea infestations, as well as the treatment and control of ear mite infestations and intestinal parasite infections in cats and kittens 9 weeks of age and older and that weigh at least 2 lbs.**

**Once-a-month topical solution for ferrets for the prevention of heartworm disease, kills adult fleas, and is indicated for the treatment of flea infestations. Indicated for ferrets that weigh at least 2 lbs.**

**CAUTION:**

Federal Law restricts this drug to use by or on the order of a licensed veterinarian.

**DESCRIPTION:**

*Advantage Multi for Cats* (10 % imidacloprid + 1 % moxidectin) is a colorless to yellow ready-to-use solution packaged in single-dose applicator tubes for topical treatment of cats. The formulation and dosage schedule are designed to provide a minimum of 4.5 mg/lb (10.0 mg/kg) imidacloprid and 0.45 mg/lb (1.0 mg/kg) moxidectin based on body weight.

Imidacloprid is a chloronicotinyl nitroguanidine insecticide. The chemical name of imidacloprid is 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine. Moxidectin is a semisynthetic macrocyclic lactone endectocide derived from the actinomycete *Streptomyces cyaneogriseus noncyanogenus*. The chemical name of moxidectin is [6R, 23E, 25S(E)]-5-0-Demethyl-28-deoxy-25-(1,3-dimethyl-1-butenyl)-6,28-epoxy-23-(methoxyimino) milbemycin B.

**INDICATIONS:**

*Advantage Multi for Cats* is indicated for the prevention of heartworm disease caused by *Dirofilaria immitis*. *Advantage Multi for Cats* kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations. *Advantage Multi for Cats* is also indicated for the treatment and control of ear mite (*Otodectes cynotis*) infestations and the following intestinal parasites:

Intestinal Parasite	Intestinal Stage		
	Adult	Immature Adult	Fourth Stage Larvae

Hookworm Species	<i>Ancylostoma tubaeforme</i>	X	X	X
Roundworm Species	<i>Toxocara cati</i>	X		X

## **WARNINGS:**

**Do not use on sick, debilitated, or underweight cats (See ADVERSE REACTIONS).**

**Do not use on cats less than 9 weeks of age or less than 2 lbs. body weight.**

## **HUMAN WARNINGS:**

**Not for human use. Keep out of the reach of children.**

**Children should not come in contact with the application site for 30 minutes after application.**

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Exposure to the product has been reported to cause headache; dizziness; and redness, burning, tingling, or numbness of the skin.

**Wash hands thoroughly with soap and warm water after handling.**

If contact with eyes occurs, hold eyelids open and flush with copious amounts of water for 15 minutes. If eye irritation develops or persists, contact a physician. If swallowed, call poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. People with known hypersensitivity to benzyl alcohol, imidacloprid or moxidectin should administer the product with caution. In case of allergic reaction, contact a physician. If contact with skin or clothing occurs, take off contaminated clothing. Wash skin immediately with plenty of soap and water. Call a poison control center or physician for treatment advice.

The Safety Data Sheet (SDS) provides additional occupational safety information. For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.

## **PRECAUTIONS:**

Do not dispense dose applicator tubes without complete safety and administration information.

Avoid oral ingestion. Cats may experience hypersalivation, tremors, vomiting and decreased appetite if *Advantage Multi for Cats* is inadvertently administered orally or through grooming/licking of the application site.

The safety of *Advantage Multi for Cats* has not been established in breeding, pregnant, or lactating cats.

The effectiveness of *Advantage Multi for Cats* against heartworm infections (*D. immitis*) after bathing has not been evaluated in cats.

Use of this product in geriatric patients with subclinical conditions has not been adequately studied. Several otherwise healthy, thin geriatric cats experienced prolonged lethargy and sleepiness after using this drug.

**(See ADVERSE REACTIONS.)**

### **ADVERSE REACTIONS:**

**Field Study:** Following treatment with *Advantage Multi for Cats* or an active control, cat owners reported the following post-treatment reactions:

<b>OBSERVATION</b>	<b>Advantage Multi n = 113</b>	<b>Active Control n = 38</b>
Lethargy (protracted sleeping, poorly responsive)	3 cats* (2.7 %)	None observed
Behavioral changes (e.g., agitated, excessive grooming, hiding, pacing, spinning)	9 cats (8.0 %)	1 cat (2.6 %)
Discomfort (e.g., scratching, rubbing, head-shaking)	5 cats (4.4 %)	None observed
Hypersalivation (within 1 hour after treatment)	3 cats (2.7 %)	None observed
Polydipsia	3 cats (2.7 %)	None observed
Coughing and gagging	1 cat (0.9 %)	None observed

\* These three cats were from the same household and included one 13-yr-old cat in good health, one 15-yr-old FIV positive cat in good health, and one 15-yr-old, underweight cat in fair health. Lethargy was noted for 24 to 36 hrs after the first treatment only; one cat was unsteady at 48 hrs. These cats were not on other medications.

During another field study, a 16-year-old cat with renal disease slept in the same place without moving for two days following application.

**(See PRECAUTIONS.)**

**Laboratory Effectiveness Studies:** *Advantage Multi for Cats* was administered at the recommended dose to 215 cats in 20 effectiveness studies. One random-sourced cat exhibited signs consistent with either moxidectin toxicity or viral respiratory disease and died 26 hours after product application; necropsy findings were inconclusive as to the cause of death. A second cat that became ill 3 days after application of *Advantage Multi for Cats* responded to treatment for respiratory infection and completed the study. A third cat became ill on day 3 and died with signs and lesions attributable to panleukopenia on day 7 after moxidectin application.

**Post-Approval Experience:** The following adverse events are based on post-approval adverse drug experience reporting. Not all adverse reactions are reported to FDA CVM. It is not always possible to reliably estimate the adverse event frequency or establish a causal relationship to product exposure using this data. The following adverse events in cats are listed in decreasing order of reporting frequency: hypersalivation, depression/lethargy, application site reactions (alopecia, pruritus, lesions, and

erythema), decreased appetite, vomiting, hyperactivity, ataxia, trembling, and behavior disorder (hiding).

In some cases death has been reported.

In humans, ocular and dermal irritation, nausea, numbness or tingling of the mouth and lips, anaphylaxis, pruritus, vomiting, and tongue/taste abnormalities have been reported following exposure to this product.

### CONTACT INFORMATION:

For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.

For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

### DOSAGE AND ADMINISTRATION:

The recommended minimum dose is 4.5 mg/lb (10.0 mg/kg) imidacloprid and 0.45 mg/lb (1.0 mg/kg) moxidectin, once a month, by topical administration.

Do not apply to irritated skin.

1. Remove one dose applicator tube from the package. As specified in the following table, administer the entire contents of the *Advantage Multi for Cats* tube that correctly corresponds with the body weight of the cat.

Cat (lbs.)	Advantage Multi For Cats	Volume (mL)	Imidacloprid (mg)	Moxidectin (mg)
2-5	<i>Advantage Multi 5</i>	0.23	23	2.3
5.1-9	<i>Advantage Multi 9</i>	0.4	40	4
9.1-18*	<i>Advantage Multi 18</i>	0.8	80	8

\* Cats over 18 lbs. should be treated with the appropriate combination of *Advantage Multi for Cats* tubes.



2. While holding the tube in an upright position, remove the cap from the tube.
3. Turn the cap over and push the other end of cap onto the tip of the tube.
4. Twist the cap to break the seal and then remove cap from the tube.



5. Part the hair on the back of the cat's neck at the base of the head, until the skin is visible. Place the tip of the tube on the skin and apply the entire contents directly on the exposed skin. Lift tube away from skin before releasing pressure on tube.

Do not get this product in the cat's mouth or eyes or allow the cat to lick the application site for 30 minutes. Treatment at the base of the head will minimize the opportunity for ingestion by grooming. In households with multiple pets, keep animals separated to prevent licking of the application site.

Stiff, matted hair or a damp, oily appearance of the hair may be observed at the application site on some cats. This is temporary and does not affect the safety and effectiveness of the product.

**Heartworm Prevention:** For prevention of heartworm disease, *Advantage Multi for Cats* should be administered at one-month intervals. *Advantage Multi for Cats* may be administered year-around or at a minimum should start one month before the first expected exposure to mosquitoes and should continue at monthly intervals until one month after the last exposure to mosquitoes. If a dose is missed and a 30-day interval between doses is exceeded, administer *Advantage Multi for Cats* immediately and resume the monthly dosing schedule. When replacing another heartworm preventative product in a heartworm prevention program, the first treatment with *Advantage Multi for Cats* should be given within one month of the last dose of the former medication. At the discretion of the veterinarian, cats older than 6 months of age may be tested to determine the presence of existing heartworm infection before treatment with *Advantage Multi for Cats* (**See ADVERSE REACTIONS - Post-Approval Experience**).

**Flea Treatment:** For the treatment of flea infestations, *Advantage Multi for Cats* should be administered at one-month intervals. If the cat is already infested with fleas when the first dose of *Advantage Multi for Cats* is administered, adult fleas on the cat will be killed. However, re-infestation from the emergence of pre-existing pupae in the environment may continue to occur for six weeks or longer after treatment is initiated. Cats treated with imidacloprid, including those with pre-existing flea allergy dermatitis have shown clinical improvement as a direct result of elimination of fleas from the cat.

**Ear Mite Treatment:** For the treatment of ear mites (*Otodectes cynotis*), *Advantage Multi for Cats* should be administered once as a single topical dose. Monthly use of *Advantage Multi for Cats* will control any subsequent ear mite infestations.

**Intestinal Nematode Treatment:** For the treatment and control of intestinal hookworm infections caused by *Ancylostoma tubaeforme* (adults, immature adults and fourth stage larvae) and roundworm infections caused by *Toxocara cati* (adults and fourth stage larvae), *Advantage Multi for Cats* should be administered once as a single topical dose.

## **ANIMAL SAFETY:**

**Studies in Kittens:** *Advantage Multi for Cats* was topically applied at 0, 1, 3, and 5X the maximum dose to 48 healthy 9-week-old kittens on days 0, 28, and 56. Lethargy was observed in 1 kitten from the 3X group and 1 from the 5X group on the day after initial treatment; the kitten from the 3X group was also disoriented and ataxic. One kitten from the 3X group had a slow pupillary light response two days after treatment and one had tremors the day after treatment. Hypersalivation was seen in one kitten from the 5X group approximately six hours post-treatment. One kitten from the 3X group was scratching at the treatment site 2 days after treatment. Slight cough was noted in 7 different kittens (2-0X, 2-1X, and 3-5X) during the 13-day period following the first treatment. Histopathology showed granulomatous inflammation at the treatment site in three 1X kittens. Causal relationship to the drug could not be determined. Pulmonary inflammation (1-5X) and lymphoid hyperplasia (2-1X, 4-3X) were seen in treated kittens. In a second study, *Advantage Multi for Cats* was topically applied at 0, 1.7, 5.2 and 8.7X the maximum dose to 48 healthy 9-week-old kittens every two weeks for 6 doses. One kitten in the 8.7X group apparently ingested an unknown amount of the drug and developed the following clinical signs prior to euthanasia: mydriasis, salivation, depression, vomiting, unsteadiness, rapid to slow to difficult breathing, poor pupillary response, generalized tremors, inability to move, and nystagmus. Two kittens in the 5.2X group developed mydriasis, salivation, depression, squinting, and poor appetite. A kitten in the 1.7X group developed mydriasis.

**Dose Tolerance Study:** Eight healthy juvenile cats were topically dosed with a single application of *Advantage Multi for Cats* at 10 times the recommended dose volume. Mild, transient hypersalivation occurred in two of the cats.

**Oral Study in Cats:** The oral safety of *Advantage Multi for Cats* was tested in case of accidental oral ingestion. The maximum topical dose was orally administered to twelve healthy 9-week-old kittens. Hypersalivation (8 of 12 kittens) and vomiting (12 of 12 kittens) were observed immediately post-treatment. Tremors developed in one kitten within 1 hour, resolving without treatment within the next hour. All 12 kittens were either anorexic or had decreased appetite for at least 1 day following treatment. In 3 kittens, the anorexia or decreased appetite continued into the second week following treatment. There was a post-treatment loss of body weight in treated kittens compared to control kittens. In a pilot safety study using kittens younger in age and lighter in weight than allowed by product labeling, an 8-week old kitten weighing 0.6 kg orally received 2X of the label topical dose (0.46 mL/kg). Immediately after dosing, it vomited, had labored breathing and slight tremors. Within 4 hours, it was normal, but was found dead on day 6. Necropsy could not determine the cause of death.

**Study in Heartworm Positive Cats:** Young adult cats were inoculated subcutaneously with third-stage *D. immitis* larvae. At 243–245 days post-infection, immunoserology and echocardiography were performed to identify cats with adult heartworm burdens similar to naturally-acquired infections. Two groups were treated topically with either *Advantage Multi for Cats* at the label dose or placebo, once every 28 days, for three consecutive treatments. A third group was treated topically, once, with *Advantage Multi for Cats* at 5X the label dose. Sporadic vomiting and labored breathing related to heartworm burden were observed in the treatment and control groups. There was no difference between treatment groups in the numbers of adult *D. immitis* recovered at study conclusion. No adverse reactions were associated with the topical application of *Advantage Multi for Cats* to experimentally heartworm-infected cats.

## **FERRETS**

**Use only the 0.4 mL ADVANTAGE MULTI for Cats in ferrets. The 0.23 mL size does not provide an effective dose and the 0.8mL size could result in an overdose.**

### **INDICATIONS:**

#### **For ferrets:**

*Advantage Multi for Cats* is indicated for the prevention of heartworm disease in ferrets caused by *Dirofilaria immitis*. *Advantage Multi for Cats* kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations on ferrets.

### **WARNINGS:**

**Do not use on sick or debilitated ferrets.**

### **PRECAUTIONS:**

Do not dispense dose applicator tubes without complete safety and administration information.

The safety of *Advantage Multi for Cats* has not been established in breeding, pregnant, and lactating ferrets.

Treatment of ferrets weighing less than 2.0 lbs (0.9 kg) should be based on a risk-benefit assessment.

The effectiveness of *Advantage Multi for Cats* in ferrets weighing over 4.4 lbs (2.0 kg) has not been established.

### **ADVERSE REACTIONS:**

**Field Safety Study in Ferrets:** *Advantage Multi for Cats* was topically administered to 131 client-owned ferrets at the recommended dose volume (0.4 mL). The ferrets ranged in age from 3 months to 7 years, and weighed between 0.5 and 1.86 kg (1.1 to 4.1 lbs). The dose of imidacloprid ranged between 21.5 to 80.2 mg/kg in this study. The dose of moxidectin ranged between 2.2 to 8.0 mg/kg in this study.

Adverse reactions in ferrets following treatment included: pruritus/scratching, scabbing, redness, wounds and inflammation at the treatment site; lethargy; and chemical odor. These adverse reactions resolved without additional therapy. Owners also reported stiffening of the hair at the treatment site, however, this is expected with application of a topical product and is not considered an adverse reaction.

Three human adverse reactions were reported. An owner's finger became red following skin contact with the product. One owner reported a headache caused by the chemical odor of the product. One owner reported a tingling sensation of the lips after kissing the treatment site.

**Foreign Market Experience:** Because the following events were reported voluntarily

during post-approval use of the product in foreign markets, it is not always possible to reliably establish a causal relationship to drug exposure. Adverse events reported in ferrets topically treated with 0.4 mL imidacloprid + moxidectin for cats included: malaise, vomiting, diarrhea, shaking, mydriasis, hypersalivation with abnormal neurologic signs, seizures, death, generalized hematoma of the body, and alopecia at the treatment site. Adverse reactions in humans included: burning, tingling, numbness, bad taste in the mouth, dizziness, and headache.

## **ANIMAL SAFETY:**

**Ferrets:** *Advantage Multi for Cats* was topically applied at 5X the recommended dose volume to six healthy 9-month-old ferrets on Study Days 0, 14, 28, and 42. Because the weights of the ferrets in this study ranged from 2.0 to 4.0 lb (0.9 kg to 1.8 kg), ferrets received a range of dosages from 51.0 to 106.9 mg/lb (112 to 235 mg/kg) of imidacloprid and 5 to 10.5 mg/lb (11 to 23 mg/kg) of moxidectin. The following abnormal clinical signs were reported during the study: wet, matted, and/or greasy appearance to the hair, shaking of the head and/or body, rubbing of dose site on cage, and shedding. Slight increases in phosphorous, potassium, aspartate aminotransferase (AST), and glucose were seen during the study, however, no clinical signs related to these bloodwork changes were reported.

**Oral Safety Study:** *Advantage Multi for Cats* was orally administered at the recommended dose volume (0.4 mL) to eight healthy ferrets on Study Day 0. Ferrets were 78 to 101 days old (11.1 to 14.4 weeks) and weighed between 1.1 to 1.8 lb (0.5 to 0.8 kg) body weight on the day of dosing, resulting in doses ranges of 22.0–36.8 mg/lb (48.3–81.0 mg/kg) imidacloprid and 2.2–3.7 mg/lb (4.8–8.0 mg/kg) moxidectin. The following abnormal clinical signs were reported immediately following oral administration of *Advantage Multi for Cats*: vomiting (one ferret) and ataxia (two ferrets). All abnormalities resolved without treatment or supportive care.

## **DOSAGE AND ADMINISTRATION:**

### **For ferrets:**

The recommended minimum dose for a ferret is 9 mg/lb (20 mg/kg) imidacloprid and 0.9 mg/lb (2 mg/kg) moxidectin, once a month, by topical administration.

<b>Ferret (lbs.)</b>	<b>Advantage Multi For Cats</b>	<b>Volume (mL)</b>	<b>Imidacloprid (mg)</b>	<b>Moxidectin (mg)</b>
2.0–4.4	<i>Advantage Multi 9</i>	0.4	40	4

**Only the 0.4 mL applicator tube volume (Advantage Multi 9) should be used on ferrets.**

Do not apply to irritated skin.

1. Remove one dose applicator tube from the package. Administer the entire contents

of the *Advantage Multi for Cats* tube (0.4 mL).



2. While holding the tube in an upright position, remove the cap from the tube.
3. Turn the cap over and push the other end of cap onto the tip of the tube.
4. Twist the cap to break the seal and then remove cap from the tube.



5. Part the hair on the back of the ferret's neck at the base of the head, until the skin is visible. Place the tip of the tube on the skin and apply the entire contents directly on the exposed skin. Lift tube away from skin before releasing pressure on tube.

Do not get this product in the ferret's mouth or eyes or allow the ferret to lick the application site for 30 minutes. Treatment at the base of the head will minimize the opportunity for ingestion by grooming. In households with multiple pets, keep animals separated to prevent licking of the application site.

Stiff, matted hair or a damp, oily appearance of the hair may be observed at the application site on some ferrets. This is temporary and does not affect the safety or effectiveness of the product.

**Heartworm Prevention:** For prevention of heartworm disease, *Advantage Multi for Cats* should be administered at one-month intervals. *Advantage Multi for Cats* may be administered year-round or at a minimum should start one month before the first expected exposure to mosquitoes and should continue at monthly intervals until one month after the last exposure to mosquitoes. If a dose is missed and a 30-day interval between doses is exceeded, administer *Advantage Multi for Cats* immediately and resume the monthly dosing schedule.

**Flea Treatment:** For the treatment of flea infestations on ferrets, *Advantage Multi for Cats* should be administered at one-month intervals. If the ferret is already infested with fleas when the first dose of *Advantage Multi for Cats* is administered, adult fleas on the ferret will be killed. However, re-infestation from the emergence of pre-existing pupae in the environment may continue to occur for six weeks or longer after treatment is initiated.

#### **STORAGE INFORMATION:**

Store at temperatures between 4 °C (39 °F) and 25 °C (77 °F), avoiding excess heat or cold.

**HOW SUPPLIED:**

Applications Per Package

3 x 0.23 mL tubes

6 x 0.4 mL tubes

6 x 0.8 mL tubes

PA513355X

Revised: August 2025

Approved by FDA under NADA # 141-254

© 2025 Elanco or its affiliates.

Advantage Multi, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

Product of Germany.

**Elanco™**

Manufactured for:

Elanco US Inc

Indianapolis, IN 46221 U.S.A.

**Principal Display Panel - 0.23 mL Carton Label**

**2 -**

**5 lbs**

Heartworm

Fleas

Hookworm

Roundworm

Ear Mites

**3 Pack**

**advantage multi™  
for cats**

**(imidacloprid+moxidectin)**

**Topical Solution**

**CAUTION: Federal Law restricts this drug to  
use by or on the order of a licensed veterinarian**

**Once-A-Month Topical Solution**

- Prevention of heartworm disease
- Kills adult fleas and is indicated for the treatment of flea infestations
- Treatment and control of ear mite infestations
- Treatment and control of hookworms and roundworms
- For cats and kittens 9 weeks of age and older and 2 -5 lbs.

**Children should not come in contact with the application site for 30 minutes after application.**

**Do not use the 0.23 mL size tube in ferrets as it does not provide an effective dose.**

Each tube contains 23 mg of imidacloprid and 2.3 mg of moxidectin.

**Keep this and all drugs out of reach of children.**

Approved by FDA under NADA # 141-254

**THREE 0.23 mL Tubes**



## Principal Display Panel - 0.4 mL Carton Label

**5.1-9 lbs**

for cats:

Heartworm

Fleas

Hookworm

Roundworm

Ear Mites

for  
ferrets:

Heartworm

Fleas

## **6 Pack**

**advantage multi™  
for cats**

**(imidacloprid+moxidectin)**

### **Topical Solution for Cats and Ferrets**

**CAUTION: Federal Law restricts this drug to  
use by or on the order of a licensed veterinarian**

### **Once-A-Month Topical Solution**

#### **For Cats:**

- Prevention of heartworm disease
- Kills adult fleas and is indicated for the treatment of flea infestations
- Treatment and control of ear mite infestations
- Treatment and control of hookworms and roundworms
- For cats and kittens 9 weeks of age and older and 5.1–9 lbs.

#### **For Ferrets:**

- Prevention of heartworm disease
- Kills adult fleas and treats flea infestations
- For ferrets that weigh at least 2 lbs.

**Children should not come in contact with  
the application site for 30 minutes after application.**

Each tube contains 40 mg of imidacloprid and 4 mg of moxidectin.

**Keep this and all drugs out of reach of children.**

Approved by FDA under NADA # 141-254

**SIX 0.4 mL Tubes**



## Principal Display Panel - 0.8 mL Carton Label

**9.1 -  
18 lbs**

Heartworm

Fleas

Hookworm

Roundworm

Ear Mites

**6 Pack**

**advantage multi™  
for cats**

**(imidacloprid+moxidectin)**

**Topical Solution**

**CAUTION: Federal Law restricts this drug to use by or on the order of a licensed veterinarian**

**Once-A-Month Topical Solution**

- Prevention of heartworm disease
- Kills adult fleas and is indicated for the treatment of flea infestations
- Treatment and control of ear mite infestations
- Treatment and control of hookworms and roundworms
- For cats and kittens 9 weeks of age and older and 9.1 -18 lbs.

**Children should not come in contact with the application site for 30 minutes after application.**

**Do not use the 0.8 mL size tube in ferrets as it could result in an overdose.**

Each tube contains 80 mg of imidacloprid and 8 mg of moxidectin.

**Keep this and all drugs out of reach of children.**

Approved by FDA under NADA # 141-254

**SIX 0.8 mL Tubes**



## ADVANTAGE MULTI FOR CATS

imidacloprid, moxidectin solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:58198-0053
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IMIDACLOPRID (UNII: 3BN7M937V8) (IMIDACLOPRID - UNII:3BN7M937V8)	IMIDACLOPRID	100 mg in 1 mL

**MOXIDECTIN** (UNII: NGU5H31YO9) (MOXIDECTIN - UNII:NGU5H31YO9) | MOXIDECTIN | 10 mg in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58198-0053-1	3 in 1 CARTON		
1	NDC:58198-0053-3	0.23 mL in 1 TUBE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141254	01/19/2007	

## ADVANTAGE MULTI FOR CATS

imidacloprid, moxidectin solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:58198-0054
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IMIDACLOPRID</b> (UNII: 3BN7M937V8) (IMIDACLOPRID - UNII:3BN7M937V8)	IMIDACLOPRID	100 mg in 1 mL
<b>MOXIDECTIN</b> (UNII: NGU5H31YO9) (MOXIDECTIN - UNII:NGU5H31YO9)	MOXIDECTIN	10 mg in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58198-0054-1	6 in 1 CARTON		
1	NDC:58198-0054-3	0.4 mL in 1 TUBE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141254	01/19/2007	

## ADVANTAGE MULTI FOR CATS

imidacloprid, moxidectin solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:58198-0055
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IMIDACLOPRID</b> (UNII: 3BN7M937V8) (IMIDACLOPRID - UNII:3BN7M937V8)	IMIDACLOPRID	100 mg in 1 mL
<b>MOXIDECTIN</b> (UNII: NGU5H31YO9) (MOXIDECTIN - UNII:NGU5H31YO9)	MOXIDECTIN	10 mg in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58198-0055-1	6 in 1 CARTON		
1	NDC:58198-0055-3	0.8 mL in 1 TUBE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141254	01/19/2007	

**Labeler** - Elanco US Inc. (966985624)

Revised: 10/2025

Elanco US Inc.