

SELENIOUS ACID- selenious acid injection, solution
Pangea Pharmaceuticals, LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

SELENIUM INJECTION
Rx Only
TRACE ELEMENT ADDITIVE FOR IV USE AFTER DILUTION

DESCRIPTION

Selenium Injection is a sterile, nonpyrogenic solution for use as an additive to solutions for Total Parenteral Nutrition (TPN). Each mL contains Selenious Acid 65.4 mcg (equivalent to elemental selenium 40 mcg/mL) and Water for Injection q.s. pH may be adjusted with nitric acid to 1.8 to 2.4.

CLINICAL PHARMACOLOGY

Selenium is part of glutathione peroxidase which protects cell components from oxidative damage due to peroxides produced in cellular metabolism.

Prolonged TPN support in humans has resulted in selenium deficiency symptoms which include muscle pain and tenderness. The symptoms have been reported to respond to supplementation of TPN solutions with selenium.

Pediatric conditions, Keshan disease, and Kwashiorkor, have been associated with low dietary intake of selenium. The conditions are endemic to geographical areas with low selenium soil content. Dietary supplementation with selenium salts has been reported to reduce the incidence of the conditions among affected children.

Normal blood levels of selenium in different human populations have been found to vary and depend on the selenium content of the food consumed. Results of surveys carried out in some countries are tabulated below:

Country	Number of Samples	SELENIUM (mcg/100mL) (a)		
		Whole Blood	Blood Cells	Plasma / Serum
Canada	254 Adults	(37.9 ± 7.8)	(23.6 ± 6.0)	(14.4 ± 2.9)
England	8 (b)	26 - 37 (32)	-	-
Guatemala & Southern USA	10 Adults 9 Children (c)	19 - 28 (22) (23 ± 5)	- (36 ± 12)	- (15 ± 5)
New		(5.4 ±	(6.6 ±	

New Zealand (d)	113 Adults	(5.4 ± 0.1)	(0.0 ± 0.3)	(4.3 ± 0.1)
Thailand	3 Adults 9 Children (e)	14.4 - 20.2 (12.0 ± 3.6) (f)	17.8 - 35.8 (19.5 ± 8.2)	8.1 - 12.5 (8.3 ± 2.2)
USA	210 Adults	15.7 - 25.6 (20.6)	-	-

(a) Mean values with or without standard deviation in parentheses, all other ranges.

(b) Age group unknown.

(c) Three children recovered from Kwashiorkor and the other six under treatment for other diseases.

(d) Low selenium-content soil area.

(e) Well nourished children, three recovered from Kwashiorkor and the other six under treatment for other diseases.

(f) Mean values from seven subjects.

Plasma selenium levels of 0.3 and 0.9 mcg/100 mL have been reported to produce deficiency symptoms in humans.

Selenium is eliminated primarily in urine. However, significant endogenous losses through feces also occur. The rate of excretion and the relative importance of two routes varies with the chemical form of selenium used in supplementation. Ancillary routes of elimination are lungs and skin.

INDICATIONS AND USAGE

Selenium Injection is indicated for use as a supplement to intravenous solutions given for total parenteral nutrition (TPN). Administration of selenium in TPN solutions helps to maintain plasma selenium levels and to prevent depletion of endogenous stores and subsequent deficiency symptoms.

CONTRAINDICATIONS

Selenium Injection should not be given undiluted by direct injection into a peripheral vein because of the potential for infusion phlebitis.

WARNING

Selenium Injection can be toxic if given in excessive amounts. Supplementation of TPN solution with selenium should be immediately discontinued if toxicity symptoms are observed. Frequent determination of plasma selenium levels during TPN support and close medical supervision is recommended.

Selenium Injection is a hypotonic solution and should be administered in admixtures only.

This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates

are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

PRECAUTIONS

As selenium is eliminated in urine and feces, selenium supplements may be adjusted, reduced or omitted in renal dysfunction and/or gastrointestinal malfunction. In patients receiving blood transfusions, contribution from such transfusions should also be considered. Frequent selenium plasma level determinations are suggested as a guideline.

In animals, selenium has been reported to enhance the action of Vitamin E and decrease the toxicity of mercury, cadmium and arsenic.

Pregnancy

Teratogenic Effects

Pregnancy Category C: Selenium at high dose levels (15-30 mcg/egg) has been reported to have adverse embryological effects among chickens. There are however, no adequate and well controlled studies in pregnant women. Selenium Injection should be used during pregnancy only if potential benefit justifies the potential risk to the fetus.

Presence of selenium in placenta and umbilical cord blood has been reported in humans.

ADVERSE REACTIONS

The amount of selenium present in Selenium Injection is small. Symptoms of toxicity from selenium are unlikely to occur at the recommended dosage level. To report SUSPECTED ADVERSE REACTIONS, contact Pangea Pharmaceuticals, LLC at 1-855-892-8224 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

OVERDOSAGE

Chronic toxicity in humans resulting from exposure to selenium in industrial environments, intake of foods grown in seleniferous soils, use of selenium-contaminated water, and application of cosmetics containing selenium has been reported in literature. Toxicity symptoms include hair loss, weakened nails, dermatitis, dental defects, gastrointestinal disorders, nervousness, mental depression, metallic taste, vomiting, and garlic odor of breath and sweat. Acute poisoning due to ingestion of large amounts of selenium compounds has resulted in death with histopathological changes including fulminating peripheral vascular collapse, internal vascular congestion, diffusely hemorrhagic, congested and edematous lungs, brick-red color gastric mucosa. The death was preceded by coma.

No effective antidote to selenium poisoning in humans is known. Animal studies have shown casein and linseed oil in feeds, reduced glutathione, arsenic, magnesium sulfate,

and bromobenzene to afford limited protection.

DOSAGE AND ADMINISTRATION

Selenium Injection provides 40 mcg selenium/mL. For metabolically stable adults receiving TPN, the suggested additive dosage level is 20 to 40 mcg selenium/day. For pediatric patients, the suggested additive dosage level is 3 mcg/kg/day.

In adults, selenium deficiency states resulting from long-term TPN support, selenium as selenomethionine or selenious acid, administered intravenously at 100 mcg/day for a period of 24 and 31 days, respectively, has been reported to reverse deficiency symptoms without toxicity.

Aseptic addition of Selenium Injection to parenteral nutrition solutions under laminar flow hood is recommended. Selenium is physically compatible with the electrolytes and other trace elements usually present in parenteral nutrition formulations. Frequent monitoring of plasma selenium levels is suggested as a guideline for subsequent administration. The normal whole blood range for selenium is approximately 10 to 37 mcg/100 mL.

Do not directly mix ascorbic acid injection with selenium containing parenteral products in the same syringe or vial, as this admixture may cause the formation of an insoluble precipitate.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

Selenium Injection containing selenious acid 65.4 mcg/mL (equivalent to elemental selenium 40 mcg/mL).

HOW SUPPLIED	NDC NUMBER
Pharmacy Bulk Package 5 x 10 mL vials per carton	81279-106-05
Pharmacy Bulk Package 10 mL Single-Use Vial	81279-106-01

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) (See USP Controlled Room Temperature).

Manufactured for: Pangea Pharmaceuticals, LLC Brielle, NJ 08730

Rev. 04/2024

PRINCIPAL DISPLAY PANEL - Vial Labeling

NDC 81279-106-01
10 mL Single Use Vial
Rx Only

Selenious Acid Injection, USP

400 mcg*/10 mL

(40 mcg*/mL) of selenium

For intravenous use after dilution and admixing

PRESERVATIVE FREE.

10 mL

PHARMACY BULK PACKAGE

<p>NDC 81279-106-01 Selenious Acid Injection, USP 400mcg*/10mL (40 mcg*/mL of Selenium) For Intravenous use after dilution and all mixing. Rx Only • Single Dose Vial 10 mL Preservative Free</p>	<p>Each mL provides: *40 mcg of selenium (present as 65.4 mcg of selenious acid).</p> <p>Recommended Dose: See prescribing information Contains no more than 2,500 mcg/L of aluminum. Store at 20° to 25°C (68° to 77° F) [See USP]. DISCARD VIAL WITHIN 4 HOURS OF ENTRY</p> <p>Manufactured for: Pangea Pharmaceuticals, LLC Rev. 04/24</p>	 <p>3 81279 10601 4 LOT: ZZZXXYY EXP: 99/9999</p>
--	---	--

PRINCIPAL DISPLAY PANEL - Carton Labeling

NDC 81279-106-05

Rx Only

Selenious Acid Injection, USP

400 mcg*/10 mL

(40 mcg*/mL) of selenium

For intravenous use after dilution and admixing

STERILE

PRESERVATIVE FREE.

5x 10 mL VIALS

PHARMACY BULK PACKAGE

NDC 81279-106-05

Rx only

Selenious Acid Injection, USP

400 mcg*/10mL

(40 mcg*/mL of Selenium)

For intravenous use after dilution and admixing.
10mL

Preservative Free

5 x 10ml Single Dose Vials

Each mL provides:
*40 mcg of selenium (present as
65.4 mcg of selenious acid)

Recommended dose:
See prescribing information

Contains no more than 2,500 mcg/L
of Aluminum.

Store at 20° to 25° C (68° to 77° F)
See USP

DISCARD VIAL WITHIN 4 HOURS OF ENTRY

Manufactured for:
Pangea Pharmaceuticals, LLC
Rev. 04/2024

GTIN: 00381279106052
SN: QWERTYUIOPAS
EXP: MM-YYYY
LOT: RRTMM



SELENIOS ACID

selenious acid injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:81279-106
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SELENIUM (UNII: H6241UJ22B) (SELENIUM - UNII:H6241UJ22B)	SELENIUM	40 ug in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
NITRIC ACID (UNII: 411VRN1TV4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81279-106-05	5 in 1 BOX	05/20/2024	
1	NDC:81279-106-01	10 mL in 1 VIAL, PHARMACY BULK PACKAGE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

unapproved drug other		05/20/2024	
--------------------------	--	------------	--

Labeler - Pangea Pharmaceuticals, LLC (117751012)

Registrant - Pangea Pharmaceuticals, LLC (117751012)

Revised: 1/2026

Pangea Pharmaceuticals, LLC