

**VICKS PAINQUIL PM LIQUICAPS PAIN RELIEVER PLUS NIGHTTIME SLEEP-AID-
acetaminophen, diphenhydramine hcl capsule, liquid filled
The Procter & Gamble Manufacturing Company**

PainQuil PM Liquicaps Pain Reliever + Nighttime Sleep-Aid

Drug Facts

Active ingredients (in each LiquiCap)

Acetaminophen 500 mg
Diphenhydramine HCl 25 mg

Purpose

Pain reliever

Nighttime sleep-aid

Use

- for the temporary relief of occasional minor aches and pains with accompanying sleeplessness.

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 3 doses (2 LiquiCaps each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- Sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.

These could be signs of a serious condition.

When using this product

- avoid alcoholic beverages
- drowsiness will occur
- do not drive a motor vehicle or operate machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only one dose (2 LiquiCaps) per day (24 hours)

children under 12 yrs

do not use

Other information

store at no greater than 25°C (77°F)

Inactive ingredients

1,4 sorbitan, D&C Red No. 33, FD&C Blue No. 1, gelatin, glycerin, mannitol, polyethylene glycol 400, polyethylene glycol 600, polyvinyl acetate phthalate, povidone, propylene glycol, sorbitol, titanium dioxide, water

Questions?

1-877-881-5813

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.

‡Maximum strength dose of active ingredients per dosing interval, only use as directed.

**DIST. BY PROCTER & GAMBLE,
CINCINNATI, OH 45202**

PRINCIPAL DISPLAY PANEL - 100 LiquiCaps™

MAX STRENGTH‡

VICKS®

PainQuil™ PM

LIQUICAPS™

PAIN RELIEVER

+ NIGHTTIME SLEEP-AID

Acetaminophen, Diphenhydramine HCl

STARTS WORKING FAST

FOR RELIEF OF:

ACHES | PAINS | HEADACHE | SORE THROAT |

+ OCCASSIONAL SLEEPLESSNESS

100 LIQUICAPS™



VICKS PAINQUIL PM LIQUICAPS PAIN RELIEVER PLUS NIGHTTIME SLEEP-AID

acetaminophen, diphenhydramine hcl capsule, liquid filled

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:84126-375 |
| Route of Administration | ORAL | | |

| Drug Facts | Active Ingredients (in each LiquiCap) | Purpose | | | | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------|---------------------|---------------------------------|------------------------|-----------------------|------------|
| | Acetaminophen 500 mg | Pain Reliever | | | | |
| | Diphenhydramine HCl 25 mg | Nighttime sleep-aid | | | | |
| Use | | | | | | |
| <ul style="list-style-type: none"> • for the temporary relief of occasional minor aches and pains with accompanying sleeplessness. | | | | | | |
| Warnings | | | | | | |
| <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 3 doses (2 LiquiCaps each) in 24 hours, which is the maximum daily amount for this product.</p> <p>• with other drugs containing acetaminophen</p> <p>• 3 or more alcoholic drinks every day while using this product</p> <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> • skin redness • hives • rash If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> | | | | | | |
| Do not use | | | | | | |
| <ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • for children under 12 years of age • with any other product containing diphenhydramine, even one used on skin • liver disease • breathing problem such as emphysema or chronic bronchitis • glaucoma • difficulty in urination due to enlargement of the prostate gland | | | | | | |
| Ask a doctor before use if you have | | | | | | |
| <ul style="list-style-type: none"> • taking the blood thinning drug warfarin • taking sedatives or tranquilizers | | | | | | |
| Stop use and ask a doctor if | | | | | | |
| <ul style="list-style-type: none"> • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • Sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. These could be signs of a serious condition. | | | | | | |
| When using this product | | | | | | |
| <ul style="list-style-type: none"> • avoid alcoholic beverages • drowsiness will occur • do not drive a motor vehicle or operate machinery | | | | | | |
| If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. | | | | | | |
| Directions | | | | | | |
| <ul style="list-style-type: none"> • take only one dose (2 LiquiCaps per day (24 hours)) | | | | | | |
| <table border="0"> <tr> <td>adults & children 12 yrs & over</td> <td>2 LiquiCaps at bedtime</td> </tr> <tr> <td>children under 12 yrs</td> <td>do not use</td> </tr> </table> | | | adults & children 12 yrs & over | 2 LiquiCaps at bedtime | children under 12 yrs | do not use |
| adults & children 12 yrs & over | 2 LiquiCaps at bedtime | | | | | |
| children under 12 yrs | do not use | | | | | |
| Other information | | | | | | |
| <ul style="list-style-type: none"> • store at no greater than 25°C (77°F) | | | | | | |
| Inactive Ingredients 1, 4 sorbitan, D&G Red No. 33, FD&C Blue No. 1, gelatin, glycerin, mastic, polyethylene glycol 400, polyethylene glycol 800, polyvinyl acetate phthalate, zovonine, propylene glycol, sorbitol, titanium dioxide, water | | | | | | |
| QUESTIONS? 1-877-881-5813 | | | | | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------------------------------------------------------------------------|---------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 500 mg |
| DIPHENHYDRAMINE HCL (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M) | DIPHENHYDRAMINE HCL | 25 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|-------------------------------------------------------|----------|
| POLYVINYL ACETATE PHTHALATE (UNII: 58QVG85GW3) | |
| POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) | |
| POLYETHYLENE GLYCOL 600 (UNII: NL4J9F21N9) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| WATER (UNII: 059QF0KO0R) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| POVIDONE (UNII: FZ989GH94E) | |
| 1,4-SORBITAN (UNII: AV0Y TZ4E6J) | |
| D&C RED NO. 33 (UNII: 9DBA0SBB0L) | |
| MANNITOL (UNII: 3OWL53L36A) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SORBITOL (UNII: 506T60A25R) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|----------|
| Color | purple | Score | no score |
| Shape | OVAL | Size | 19mm |
| Flavor | | Imprint Code | PQPM |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|----------------------------------------------------|----------------------|--------------------|
| 1 | NDC:84126-375-01 | 1 in 1 CARTON | 03/17/2026 | |
| 1 | | 100 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| OTC Monograph Drug | M013 | 03/17/2026 | |

Revised: 2/2026

The Procter & Gamble Manufacturing Company