

FEVIA DIARRHEA CONTROL- bismuth subsalicylate capsule, liquid filled
VIVUNT PHARMA LLC

FEVIA Diarrhea control 48

Drug Facts

Active ingredient (in each softgel)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

Relieves:

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behaviour with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
 - 2 softgels every 1/2 hour or 4 softgels every hour as needed for diarrhea
 - 2 softgels every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 16 softgels in 24 hours
- use until diarrhea stops but no more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- store at room temperature 20°-25°C (68°-77°F)
- Protect from light, heat and moisture

Inactive ingredients

Beeswax, Colloidal Silicon Dioxide, Gelatin, Glycerin, Medium Chain Triglycerides, Polysorb, Povidone K-30, Purified Water, Sodium Carboxymethylcellulose, Sorbitol 70%, Soy Lecithin, Titanium Dioxide.

Questions?

Visit www.fevia.live

Distributed by

VIVUNT

Distributed by: VIVUNT PHARMA LLC

8950 SW 74th Court, Suite 1901

Miami, FL 33156-3178, USA.

Product from India.

www.vivunt.live

Fevia™

DIARRHEA CONTROL

Antidiarrheal

Bismuth Subsalicylate 262 mg

48 SOFTGELS

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Sugar Free

Actual size



Fevia™

DIARRHEA CONTROL

Antidiarrheal
Bismuth Subsalicylate 262 mg

48 SOFTGEL

Drug Facts

Active ingredient (in each softgel) Purpose
Bismuth subsalicylate 262 mg Upset stomach reliever and antidiarrheal

Uses
Relieves: ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin) ■ taking other salicylate products.
Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool.
Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis.
When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.
Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions
■ swallow with water, do not chew ■ adults and children 12 years and over: ■ 2 softgels every 1/2 hour or 4 softgels every hour as needed for diarrhea ■ 2 softgels every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 16 softgels in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

Other information ■ store at room temperature 20°-25°C (68°-77°F) ■ Protect from light, heat and moisture.

Inactive ingredients Beeswax, Colloidal Silicon Dioxide, Gelatin, Glycerin, Medium Chain Triglycerides, Polysorb Povidone K-30, Purified Water, Sodium Carboxymethylcellulose, Sorbitol 70%, Soy Lecithin, Titanium Dioxide.

Questions? Visit www.fevia.live

NDCXXXXXXXXXXXXXXXXXXXX

VIVUNT 30 YEARS

Distributed by: VIVUNT PHARMA LLC
8950 SW 74th Court, Suite 1901
Miami, FL 33156-3178, USA.
Product from India.
www.vivunt.live

bismuth subsalicylate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82706-028
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SOYBEAN LECITHIN (UNII: 1DI56QDM62)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POVIDONE K30 (UNII: U725QWY32X)	
BEESWAX (UNII: 2Z A36H0S2V)	
GELATIN (UNII: 2G86QN327L)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
SORBITOL SOLUTION 70% (UNII: 8KW3E207O2)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	pink (Opaque)	Score	no score
Shape	CAPSULE (oblong)	Size	12mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82706-028-01	1 in 1 CARTON	02/18/2026	
1		48 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/18/2026	

Labeler - VIVUNT PHARMA LLC (045829437)

Establishment

Name	Address	ID/FEI	Business Operations
MEDGEL PRIVATE LIMITED		677385498	manufacture(82706-028)

Revised: 2/2026

VIVUNT PHARMA LLC