GOOD NEIGHBOR PHARMACY COUGH DM ER- dextromethorphan polistirex suspension Amerisource Bergen

Amerisource Bergen Cough DM ER Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake bottle well before use
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24
	hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24
	hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24
	hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

Inactive Ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Delsym® active ingredient

12 HOUR COUGH RELIEF

Cough

DM ER

dextromethorphan polistirex

extended-release oral suspension

Cough Suppressant

Orange-Flavored Liquid

Dosing Cup Included

Alcohol-Free – Day or Night

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

3 fl oz (89 mL)





Drug Facts

Active ingredient (in each 5 mL)
Dextromethorphan polistirex equivalent to
30 mg dextromethorphan hydrobromide......

Purpose

USES temporarily relieves

■ cough due to minor threat and bronchial irritation as may occur with the common cold or hisheld irritants

■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine coldsse inhibitor
MADO) (cartain drugs for depression, psychiatric or emotional conditions, or
"Arkinson's disease), or for 2 weeks after stopping the MACI drug. If you do not
one will your prescription drug cordiains an MADI, ask a doctor or pharmacist
softere taking this product.

Malergy Alert: Contains sodium metabisuffite, a suffite that may cause

allergic-type reactions.

allargic-type reactions.

Ask a doctor before use if you have

chronic cough that lasts as occurs with snoking, asthma or emphysema

cough that occurs with too much phisign (mucus)

Stop use and ask a doctor if

side effects occur. You may report side effects to FDA at 1-800-FDA-1088.

cough lasts more than 7 days, cough comes back, or occurs with feest, rash or headsche that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In cose of overdose, get medical help or contact a Poison Centrol Center right away (1-800-222-1222).

- Directions
 shake bottle well before use
 measure only with doeing cup provided
 measure doeing cup with other products
 does as follows or as directed by a doctor
- adults and children 10 mL every 12 hours, not to exceed 20 mL in 24 hours 12 years of age and over

5 mL every 12 hours, not to exceed 10 mL in 24 hours children 6 to under 12 years of age 2.5 mL every 12 hours, not to exceed 5 mL in 24 hours children under 4 years of age do not use

Other information
■ each 5 mL contains: sodium 5 mg
■ store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

Drug Facts (continued)

Drag in active ingredients D&C Red #30 aluminum lake, D&C Vellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben natural and artificial orange flavor, polysorbate 80 polyvinyl acstate, povidone, propylparaben, purified water, sodium metabisulfile, sodium olystyrene sulfonate, sucrose, tartaric acid, ragacanth gum, triacetin, xanthan gum

Questions or comments? 1-800-719-9260

*Good Neighbor Pharmacy® Cough DM ER is not manufactured or distributed by Reckitt Benckiser,

DOSING		
SHAKE WELL BEFORE USE.		

	with dosing cup provided. ing cup with other products.
Age (yr)	Dose
12 years	10 mL
to adult	EVERY 12 HOURS
6 to	5 mL
under 12	EVERY 12 HOURS
4 to	2.5 mL
under 6	EVERY 12 HOURS





PHARMACY®

NEIGHBOR

GOOD

COUGH RELIEF

dextromethorphan polistirex extended-release oral suspension

Cough Suppressant

Orange-Flavored Liquid

Dosing Cup Included
Alcohol-Free • Day or Night

3 fl oz (89 mL



NDC 48122-141-21

► 12 HOUR







GOOD NEIGHBOR PHARMACY COUGH DM ER

dextromethorphan polistirex suspension

Product Information

HUMAN OTC DRUG NDC:46122-141 Product Type Item Code (Source)

ORAL **Route of Administration**

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength DEXTROMETHO RPHAN HYDRO BRO MIDE (UNII: 9 D2RTI9 KYH) DEXTROMETHORPHAN 30 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)

HYDROBROMIDE

in 5 mL

Inactive Ingredients

Ingredient Name

Strength

POLISTIREX (UNII: 5H9 W9 GTW27)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
METHYLPARABEN (UNII: A2I8 C7HI9 T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	
PO VIDO NE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)	
SUCROSE (UNII: C151H8M554)	
TARTARIC ACID (UNII: W4888I119H)	
TRAGACANTH (UNII: 2944357O2O)	
TRIACETIN (UNII: XHX3C3X673)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-141-21	1 in 1 CARTON	08/30/2012	
1		89 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:46122-141-25	1 in 1 CARTON	08/02/2013	
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	08/30/2012	

Labeler - Amerisource Bergen (007914906)

Revised: 7/2020 Amerisource Bergen