

ASPIRIN LOW DOSE CHEWABLE ORANGE- aspirin tablet, chewable

Major Pharmaceuticals

Drug Facts

Active ingredient (in each tablet)

Aspirin 81mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for Chewable Aspirin 81 mg

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever get worse or lasts more than 3 days
 - redness or swelling is present
 - ringing in the ears or loss of hearing occurs
 - any new symptoms occur

These could be signs of a serious condition

If pregnant or breast-feeding

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. (800-222-1222).

Directions

- drink a full glass of water with each dose
-

adults and children 12 years and over

take 4 to 8 tablets every 4 hours not exceed 48 tablets in 24 hours unless directed by a doctor

children under 12 years

consult a doctor

Other information

- store at room temperature (15⁰-30⁰C)
- **Tamper Evident:** do not use if safety seal under cap is broken or missing

Inactive ingredients

corn starch, dextrose monohydrate, FD&C yellow #6, orange flavor, silicon dioxide, sodium saccharine, stearic acid

Questions or comments?

(866) 562-2756 (Mon-Fri 8 AM to 4 PM EST)

.

Cartons of 300 chewable tablets (10 chewable tablets each blister pack x 30), NDC 0904-7509-30

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. Keep this and all drugs out of the reach of children.

Packaged and Distributed by:

MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268 USA

Refer to package label for Distributor's NDC Number

Rev. 12/23	Re-order No. 701003	M-111
------------	---------------------	-------

Principal Display Panel

MAJOR®

0904-7509-30

Unit Dose

Aspirin Low Dose Chewable Orange

Pain Reliever (NSAID)

Tablets

81 mg

300 TABLETS (30 x 10)

Keep this and all drugs out of the reach of children. This Unit Dose package is not child resistant

and is Intended for Institutional use Only.

The drug product contained in this package is from NDC # 0904-6794, Major® Pharmaceuticals

Packaged and Distributed by:

MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268 USA

www.major-rugby.com

Drug Facts					
Active Ingredient (in each tablet)	Purpose				
Aspirin 81 mg (NSAID) *nonsteroidal anti-inflammatory drug	Pain reliever				
Uses					
<ul style="list-style-type: none"> for the temporary relief of minor aches and pains or as recommended by your doctor ask your doctor about other uses for Chewable Aspirin 81 mg 					
Warnings					
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include <ul style="list-style-type: none"> hives facial swelling asthma (wheezing) shock </p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed </p> <p>Do not use if you are allergic to aspirin or any other pain reliever/fever reducer or for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor</p> <p>Ask a doctor before use if <ul style="list-style-type: none"> stomach bleeding warning applies to you you have a history of stomach problems such as heartburn you have high blood pressure, heart disease, liver cirrhosis or kidney disease you have asthma you are taking a diuretic </p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for <ul style="list-style-type: none"> gout diabetes arthritis </p> <p>Stop use and ask a doctor if <ul style="list-style-type: none"> an allergic reaction occurs. Seek medical help right away you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days fever get worse or lasts more than 3 days redness or swelling is present ringing in the ears or loss of hearing occurs any new symptoms occur These could be signs of a serious condition </p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222).</p>					
Directions					
<ul style="list-style-type: none"> drink a full glass of water with each dose <table border="1"> <tbody> <tr> <td>adults and children 12 years and over</td> <td>take 4 to 8 tablets every 4 hours not exceed 48 tablets in 24 hours unless directed by a doctor</td> </tr> <tr> <td>children under 12 years</td> <td>consult a doctor</td> </tr> </tbody> </table>		adults and children 12 years and over	take 4 to 8 tablets every 4 hours not exceed 48 tablets in 24 hours unless directed by a doctor	children under 12 years	consult a doctor
adults and children 12 years and over	take 4 to 8 tablets every 4 hours not exceed 48 tablets in 24 hours unless directed by a doctor				
children under 12 years	consult a doctor				
Other information					
<ul style="list-style-type: none"> store at room temperature (15°-30°C) Tamper Evident: do not use if safety seal under cap is broken or missing 					
Inactive ingredients corn starch, dextrose monohydrate, FD&C yellow #6, orange flavor, silicon dioxide, sodium saccharine, stearic acid					
Questions or comments? (866) 962-2766 (Mon-Fri 8 AM to 4 PM EST)					

LBLN0294-01

Rev. 03/25

MAJOR®

NDC 0904-7509-30

Unit Dose

Aspirin Low Dose Chewable Orange Pain Reliever (NSAID)

Tablets

81 mg**300 TABLETS (30 x 10)**

Keep this and all drugs out of the reach of children. This Unit Dose package is not child resistant and is intended for Institutional Use Only.

The drug product contained in this package is from NDC # 0904-6794, Major® Pharmaceuticals

Packaged and Distributed by:
MAJOR® PHARMACEUTICALS
 Indianapolis, IN 46268 USA
www.major-rugby.com

**ASPIRIN LOW DOSE CHEWABLE ORANGE**

aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-7509(NDC:0904-6794)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	PH;034
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-7509-30	300 in 1 CARTON	05/22/2025	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/22/2025	

Labeler - Major Pharmaceuticals (191427277)