

MUCINEX FAST-MAX RECHARGE NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, triprolidine hydrochloride solution
RB Health (US) LLC

Mucinex® Nightshift
Cold and Flu

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Tripolidine HCl 2.5 mg	Antihistamine

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings Section:

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses (80 mL) in 24 hours,, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

Inactive ingredients

ammonium glycyrrhizate,
anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C red no. 40, flavor, glycerin (soy), propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 180 mL Bottle Label

NDC 72854-167-66

Mucinex®

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Triprolidine HCl – Antihistamine

6 FL OZ (180 mL)
FOR AGES 12+

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Tamper evident: Do not use if neckband on bottle cap is broken or missing.
*Helps to relieve these symptoms at night

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Maximum Strength per 4-hour dose

Dist. by: RB Health (US)
Parapet, NJ 07054-0224
©2012 RB Health
091625

LOT: 3340906
EXP.:
MADE IN:

Drug Facts

Active Ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....Pain reliever/fever reducer	
Dextromethorphan HBr 20 mg.....Cough suppressant	
Triprolidine HCl 2.5 mg.....Antihistamine	

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses (80 mL) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

PEEL HERE

Drug Facts (continued)

- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C red no. 40, flavor, glycerin (soy), propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions?
1-866-MUCINEX (1-866-682-4639)

Scan for FAQs and instructions on proper disposal of medicines or visit www.mucinex.com/FAQs

reckitt

Please visit our website
www.mucinex.com
Patents:
www.reckitt.com/patents

MUCINEX FAST-MAX RECHARGE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, triprolidine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-167
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 20 mL

Inactive Ingredients	
Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
AMMONIUM GLCYRRHIZATE (UNII: 3VRD35U26C)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	orange	Score	
Shape		Size	
Flavor	HONEY	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-167-66	180 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	03/01/2026	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2026	

Labeler - RB Health (US) LLC (081049410)

