

**EFFERVESCENT ORANGE COLD RELIEF- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent
CVS Pharmacy, Inc.**

CVS Health Effervescent Cold Relief Orange Flavor Tablets 20 ct.

Active Ingredients

Aspirin 325 mg (NSAID)*

Chlorpheniramine maleate 2 mg

Phenylephrine Bitartrate 7.8 mg

*Nonsteroidal anti-inflammatory drug

Purpose

Aspirin 325 mg (NSAID)*.Pain reliever/fever reducer

Chlorpheniramine maleate 2 mgAntihistamine

Phenylephrine Bitartrate 7.8 mgNasal decongestant

*Nonsteroidal anti-inflammatory drug

Uses

temporarily relieves these symptoms of the common cold:

- fever ■ sneezing ■ nasal congestion ■ sore throat ■ headache
- minor aches and pains ■ runny nose
- sinus congestion and pressure

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than

directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you are taking sedatives or tranquilizers
- You have
- glaucoma
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug.
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

- do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.
- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

If pregnant or breastfeeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children

Keep out of reach of children.

In case of overdose,

get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 ounces of water every 4-6 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

each tablet contains:

sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

Inactive Ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Distributed by:

CVS Pharmacy, Inc.

One CVS Drive, Woonsocket RI 02895

CVS.com

1-800 SHOP CVS

Package Label.Principle Display Panel



Drug Facts (continued)

Some throat warnings: If sore throat is severe, headache, rash, nausea, or vomiting, accompanied or followed by fever, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/reducer
- if you are allergic to aspirin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI.

Ask a doctor before use if: stomach bleeding warning applies to you; ask a doctor or pharmacist before taking this product.

- you have a history of stomach problems, such as heartburn
- you are taking blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have high blood pressure
- you are taking a diuretic
- you have glaucoma
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if: you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs.

Drug Facts

Active Ingredients (in each effervescent tablet)

Purpose

Nonsteroidal anti-inflammatory drug

Phenylephrine bitartrate 7.5 mg.....Nasal decongestant

Chlorpheniramine maleate 2 mg.....Antihistamine

Aspirin 325 mg (NSAID)*.....Pain reliever/fever reducer

Uses

temporarily relieves these symptoms of the common cold:

- fever
- sneezing
- nasal congestion
- sore throat
- headache
- minor aches and pains
- runny nose

- sinus congestion and pressure

Warnings

Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you are age 60 or older. Have had stomach ulcers or bleeding problems. Take a blood thinning (anticoagulant) or steroid drug. Take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product. Take more or for a longer time than directed.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, facial swelling

Stomach bleeding warning: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.



Orange Flavor

Effervescent Cold Relief



Actual Size



Compare to the active ingredients in Alka-Seltzer Plus® Cold Orange Zest†

Orange Flavor

NDC 69842-875-20

Effervescent Cold Relief

ASPIRIN (NSAID) - Pain reliever; Fever reducer
CHLORPHENIRAMINE MALEATE - Antihistamine
PHENYLEPHRINE BITARTRATE
 Nasal decongestant

Fast relief of:

Nasal congestion; Runny nose; Sneezing; Headache; Body ache; Sore throat; Sinus pain & pressure



Orange Flavor



Actual Size Tablet on Top Panel

20 TABLETS



Drug Facts (continued)

Seek medical help right away.

- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

Other information

each tablet contains:

- sodium 464 mg
- phenyleturonics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2017 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS
 V-17432




†This product is not manufactured, distributed or licensed by the maker of Alka-Seltzer Plus® Cold Orange Zest®.

#182122



DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

PN: 87560 CH C1
REV: 02/17



EFFERVESCENT ORANGE COLD RELIEF

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-875
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASPARTAME (UNII: Z0H242BBR1)	
DOCUSATE SODIUM (UNII: F05Q2T2JA0)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MANNITOL (UNII: 3OWL53L36A)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	ORANGE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-875-20	10 in 1 CARTON	03/10/2017	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/10/2017	

Labeler - CVS Pharmacy, Inc. (062312574)

Revised: 11/2024

CVS Pharmacy, Inc.