

LIDOCAINE PAIN RELIEF GEL -PATCH- lidocaine patch
Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.

83602-185

Active ingredient

Lidocaine 4% Purpose: Topical Anesthetic

Purpose

Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Ask a doctor if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

Stop use if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

Pregnancy or breast feeding

If pregnant or breastfeeding, ask a health professional before use

Keep out of reach of children

Keep out of reach of children. If swallowed accidentally , get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under age of 12:Consult a doctor

Other information

Store at 20°C to 25°C (68°F - 77°F)

Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, PropyleneGlycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments?

supports@povihomeofficial.com

Package label. Principal display panel

CUT OPEN HERE

Povihome®
The more attention to health. Child Resistant Packaging (Before Opening)

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

Odor-Free

Single-Use Application

Numbs Pain Away

1 PATCH
7.78" X 4.72"
20cm X 12cm

XL PATCH
Lidocaine 4%



Usage

- Remove patch from outer package
- Partially peel back protective film
- Place on affected area
- Remove remaining film while pressing patch to skin

<p>Drug Facts</p> <p>Active Ingredient Purpose Lidocaine 4% Topical anesthetic</p> <p>Uses Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.</p> <p>Warnings For external use only</p> <p>Do not use</p> <ul style="list-style-type: none"> more than 1 patch on your body at a time or with other topical anesthetics at the same time on wounds or damaged skin with a heating pad or device with other ointments, creams, sprays, or liniments if you are allergic to any ingredients of this product <p>When using this product</p> <ul style="list-style-type: none"> use only as directed avoid contact with the eyes, mucous membranes, or rashes do not bandage tightly dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch. <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering condition worsens symptoms persist for more than 7 days symptoms clear up and occur again within a few days <p>6102482017160</p>	<p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.</p> <p>Directions Adults and Children above 12 years old:</p> <ul style="list-style-type: none"> Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Apply 1 patch at a time to affected area, not more than 3 to 4 times daily. Remove patch from the skin after at most 8-hour application <p>Children under age of 12: Consult a doctor</p> <p>Other Information</p> <ul style="list-style-type: none"> Store at 20°C to 25°C (68°F - 77°F) Store in a cool dry place away from direct sunlight <p>Inactive ingredients Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, PropyleneGlycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water</p> <p>Questions or Comments: supports@povihomeofficial.com</p> <p>Child Resistant Packaging</p>
---	--

LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomeofficial.com



Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

- Odor-Free
- Single-Use Application
- Numbs Pain Away



11 PATCHES
7.87" X 4.72"
20cm X 12cm

XL PATCH
Lidocaine 4%

Guangzhou Liucuan Import & Export Trade Co., Ltd.
411 Building A, No. 8, Dong Heng San Road,
Zhucau, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

Povihome®
The more attention to health

Drug Facts

Active ingredient Lidocaine 4% **Purpose** Topical anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or lotions
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or robes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

- Adults and Children above 12 years old:
- Clean and dry the affected area.
 - Tear off the protective film and apply the exposed part of the patch to the affected area.
 - Carefully remove remaining film while pressing the patch firmly on the skin.
 - Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
 - Remove patch from the skin after at most 8-hour application.
- Children under age of 12: Consult a doctor

Other Information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive Ingredients

Dihydroxyaluminum aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone 900, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:

support@povihomeofficial.com

Child Resistant Packaging



0 542428 010337

LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomeofficial.com



Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

Odor-Free

Single-Use Application

Numbs Pain Away



12 PATCHES
7.87" X 4.72"
20cm X 12cm

XL PATCH
Lidocaine 4%

Guangzhou Liguanniao Import & Export Trade Co., Ltd.
417, Building A, No. 8, Dong Heng San Road,
Zhucaun, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

The more attention to health
Povihome®

Drug Facts

Active ingredient **Purpose**
Lidocaine 4% Topical anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days



8 9 4 2 4 8 8 0 1 5 9 0 6

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

- Adults and Children above 12 years old:**
- Clean and dry the affected area.
 - Peel off the protective film and apply the exposed part of the patch to the affected area.
 - Carefully remove remaining film while pressing the patch firmly on the skin.
 - Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
 - Remove patch from the skin after at most 8-hour application
- Children under age of 12:** Consult a doctor

Other Information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:

support@povihomeofficial.com

Child Resistant Packaging

Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

Odor-Free

Single-Use Application

Numbs Pain Away

17

PATCHES
7.87" X 4.72"
20cm X 12cm

XL PATCH

Lidocaine 4%

Guangzhou Lidocaine Import & Export Trade Co., Ltd.
472 Building A, No. 8, Dong Hong San Road,
Zhucon, Tianhe District, Guangzhou, Guangdong, China.

Usage

- 1**

Remove patch from outer package
- 2**

Partially peel back protective film
- 3**

Place on affected area
- 4**

Remove remaining film while pressing patch to skin

The more attention to health

<p>Drug Facts</p> <p>Active ingredient Purpose Lidocaine 4% Topical anesthetic</p> <p>Uses Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hip, shoulder, elbow.</p> <p>Warnings For external use only</p> <p>Do not use</p> <ul style="list-style-type: none"> • more than 1 patch on your body at a time or with other topical anesthetics at the same time • on wounds or damaged skin • with a heating pad or device • with other ointments, creams, sprays, or lotions • if you are allergic to any ingredients of this product <p>When using this product</p> <ul style="list-style-type: none"> • use only as directed • avoid contact with the eyes, mucous membranes, or rash • do not bandage tightly • dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch. <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering • condition worsens • symptoms persist for more than 7 days • symptoms clear up and occur again within a few days 	<p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.</p> <p>Directions Adults and Children above 12 years old:</p> <ul style="list-style-type: none"> • Clean and dry the affected area. • Tear off the protective film and apply the exposed part of the patch to the affected area. • Carefully remove remaining film while pressing the patch firmly on the skin. • Apply 1 patch up to 4 times to affected area, not more than 3 to 4 times daily. • Remove patch from the skin after at most 8-hour application • Children under age of 12: Consult a doctor <p>Other Information</p> <ul style="list-style-type: none"> • Store at 20°C to 25°C (68°F - 77°F) • Store in a cool dry place away from direct sunlight <p>Inactive ingredients Dihydroxyethylamine Ammonioacetate, Edetate Disodium, Glycine, Kaolin, Methylparaben, Polyacrylate 80, Propylene Glycol, Polyacrylate Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water</p> <p>Questions or Comments: support@povihomeofficial.com</p> <p>Child Resistant Packaging</p>
--	--

6 90 4 2 4 6 8 0 1 5 0 4 4

LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomemedical.com



Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

- Odor-Free
- Single-Use Application
- Numbs Pain Away



18 PATCHES
7.87" X 4.72"
20cm X 12cm

XL PATCH
Lidocaine 4%

Guangzhou Lidozhen Medical Supply & Export Trade Co., Ltd.
401 Building A, No. 8 Dong Heng San Road,
Zhuou, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

The more attention to health
Povihome®

Drug Facts

Active ingredient Lidocaine 4% **Purpose** Topical anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or mouth
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

- Adults and Children above 12 years old
 - Clean and dry the affected area.
 - Tear off the protective film and apply the exposed part of the patch to the affected area.
 - Carefully remove remaining film while pressing the patch firmly on the skin.
 - Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
 - Remove patch from the skin after at most 8-hour application.
- Children under age of 12: Consult a doctor

Other Information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyacetone, Acetaminophen, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polyorbate 80, Propylene Glycol, Polyacrylic Acid, Polysorbate 80, Polysorbate 20, Propylparaben, Sodium Polyoxyethylene Sulfate, Tartaric Acid, Water

Questions or Comments:

support@povihomemedical.com

Child Resistant Packaging



6 9 4 2 4 8 0 1 5 9 1 3

LIDOCAINE
PAIN RELIEF GEL-PATCH
www.povihomemedical.com



Povihome®
The more attention to health

Maximum OTC Strength

LIDOCAINE

PAIN RELIEF GEL-PATCH

Desensitizes Aggravated Nerves And Relieves Pain.

- Odor-Free
- Single-Use Application
- Numbs Pain Away



23 (PATCHES)
7.87" X 4.72"
20cm X 12cm

XL PATCH
Lidocaine 4%

Guangzhou Lianmeier Import & Export Trade Co., Ltd.
417 Building A, No. 8, Dongfeng Sea Road,
Zhucon, Tianhe District, Guangzhou/Guangdong, China.

Made in P.R.C.

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

Drug Facts

Active Ingredient Purpose
Lidocaine 4% Topical anesthetic

Uses
Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings
For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other treatments, creams, sprays, or lotions
- if you are allergic to any ingredients of the product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or sores
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests the patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blisters
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

The more attention to health
Povihome®

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 2 to 4 times daily.
- Remove patch from the skin after at most 8-hour application.
- Children under age of 12: Consult a doctor

Other Information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive Ingredients


Dihydroxyethylamine Acrylate, Edetate Disodium, Glycerin, Kollin, Methylparaben, Polyacrylate Gels, Polyethylene Glycol, Polycrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Terephthalic Acid, Water

Questions or Comments:

support@povihomemedical.com

Child Resistant Packaging





LIDOCAINE PAIN RELIEF GEL -PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83602-185
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.04 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

GLYCERIN (UNII: PDC6A3C0OX)
POVIDONE K90 (UNII: RDH86HJV5Z)
METHYLPARABEN (UNII: A2I8C7HI9T)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
EDETATE DISODIUM (UNII: 7FLD91C86K)
KAOLIN (UNII: 24H4NWX5CO)
TARTARIC ACID (UNII: W4888I119H)
WATER (UNII: 059QF0K00R)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)
SODIUM POLYACRYLATE (250000 MW) (UNII: 05I15JNI2J)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83602-185-11	11 in 1 BOX	02/28/2026	
1		1 in 1 BAG		
1		21 g in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:83602-185-12	12 in 1 BOX	02/28/2026	
2		1 in 1 BAG		
2		21 g in 1 PATCH; Type 0: Not a Combination Product		
3	NDC:83602-185-17	17 in 1 BOX	02/28/2026	
3		1 in 1 BAG		
3		21 g in 1 PATCH; Type 0: Not a Combination Product		
4	NDC:83602-185-18	18 in 1 BOX	02/28/2026	
4		1 in 1 BAG		
4		21 g in 1 PATCH; Type 0: Not a Combination Product		
5	NDC:83602-185-23	23 in 1 BOX	02/28/2026	
5		1 in 1 BAG		
5		21 g in 1 PATCH; Type 0: Not a Combination Product		
6	NDC:83602-185-24	24 in 1 BOX	02/28/2026	
6		1 in 1 BAG		

6

21 g in 1 PATCH; Type 0: Not a Combination Product

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/28/2026	

Labeler - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Registrant - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Revised: 2/2026

Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.