

BACTOSHIELD CHG- chlorhexidine gluconate solution
SC Johnson Professional USA, Inc.

Bactoshield CHG

Drug Facts

Active ingredient

Chlorhexidine gluconate 4% solution

Purpose

Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **skin wound and general skin cleansing**

Warnings

For external use only

Allergy alert

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient are allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.

- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Skin wound and general skin cleansing

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol.

Questions or comments?

call **1-800-587-3721** Monday through Friday 8:00 AM to 4:30 PM CST

PRINCIPAL DISPLAY PANEL - 946 mL Bottle Label

SCJ PROFESSIONAL
HEALTHCARE

Bactoshield®
CHG 4%

NDC 11084-039-24

Surgical Scrub
Chlorhexidine Gluconate 4%
Solution Antiseptic

Surgical Scrub

4% CHG


Antimicrobial

Net Contents: 32 fl oz (946 mL)

REORDER #
1344-24

1344-21K(622)
SAP # 4000011063
4SCJ32BTLLBL

deb
SKIN CARE

 <p>Bactoshield® CHG 4%</p> <p>NDC 11084-039-24</p> <p>Surgical Scrub Chlorhexidine Gluconate 4% Solution Antiseptic</p> <p>Surgical Scrub</p> <p>4% CHG</p> <p>Antimicrobial</p> <p>Net Contents: 32 fl oz (946 mL)</p> <p>REORDER # 1344-24</p> <p>1344-21K(622) SAP # 4000011063 4SCJ32BTLLBL</p> <p>deb SKIN CARE</p>	Drug Facts <p>Active ingredient Chlorhexidine gluconate 4% solution Antiseptic</p> <p>Purpose Antiseptic</p> <p>Uses</p> <ul style="list-style-type: none">■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease■ skin wound and general skin cleansing <p>Warnings For external use only</p> <p>Allergy alert: This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none">■ wheezing/difficulty breathing■ shock■ facial swelling■ hives■ rash <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none">■ if you or the patient are allergic to chlorhexidine gluconate or any other ingredient in this product■ in contact with meningitis■ in the genital area <p>When using this product</p> <ul style="list-style-type: none">■ keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.	Drug Facts (continued) <ul style="list-style-type: none">■ If solution should contact these areas, rinse out promptly and thoroughly with water■ wounds which involve more than the superficial layers of the skin should not be routinely treated■ repeated general skin cleansing of large body areas should not be done except when advised by a health care provider <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none">■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.Surgical hand scrub:<ul style="list-style-type: none">■ wet hands and forearms with water■ scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces■ a separate nail cleaner may be used■ rinse thoroughly■ wash for an additional 3 minutes with 5 mL of product and rinse under running water■ dry thoroughly <p>Healthcare personnel handwash:</p> <ul style="list-style-type: none">■ wet hands with water■ dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds■ rinse and dry thoroughly	Drug Facts (continued) <p>Skin wound and general skin cleansing:</p> <ul style="list-style-type: none">■ thoroughly rinse the area to be cleaned with water■ apply the minimum amount of product necessary to cover the skin or wound area and wash gently■ rinse again thoroughly <p>Other information</p> <ul style="list-style-type: none">■ store at 20-25°C (68-77°F)■ avoid excessive heat above 40°C (104°F) <p>Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, triethyl alcohol.</p> <p>Questions or comments? call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST</p> <p>Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.</p> <p>Manufactured for: SCJ Professional USA, Inc. Charlotte, NC 28217 1-866-789-0422 www.scjp.com</p> <p>WARNING: Cancer – www.P65Warnings.ca.gov. Keep from freezing. Made in U.S.A</p>
---	---	--	--

BACTOSHIELD CHG

chlorhexidine gluconate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11084-039
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	40 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
COCO DIETHANOLAMIDE (UNII: 92005F972D)	
GLUCONOLACTONE (UNII: WQ29KQ9POT)	
HYDROXYETHYL CELLULOSE (5000 MPA.S AT 1%) (UNII: X70SE62ZAR)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LAURAMINE OXIDE (UNII: 4F6FC4MI8W)	
PEG-75 LANOLIN (UNII: 09179OX7TB)	
WATER (UNII: 059QF0KO0R)	
TRIDECYL ALCOHOL (UNII: 8I9428H868)	

Product Characteristics

Color	YELLOW	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11084-039-03	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2022	
2	NDC:11084-039-24	946 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019125	12/01/2022	

Labeler - SC Johnson Professional USA, Inc. (607378015)

Establishment

Name	Address	ID/FEI	Business Operations
Xttrium Laboratories, Inc.		007470579	MANUFACTURE(11084-039)

Revised: 11/2022

SC Johnson Professional USA, Inc.