

DOLODOL- acetaminophen, aspirin, caffeine tablet
OPMX LLC

Dolodol EXH 20-4 w/80 caplets

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg
Aspirin 250 mg (NSAID)*
Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID)*.....Pain reliever
Caffeine 65 mg.....Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- sinusitis
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- facial swelling

- asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and, occasionally, rapid heartbeat.

Do not use

- if you ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- an allergic reaction occurs.

- Seek medical help right away
- any new symptoms appear
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- painful area is red or swollen

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over
- take 2 tablets every 6 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15-30°C (59-86°F)
- TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAPS IS BROKEN OR MISSING

Inactive ingredients

corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

Exclusively distributed by:

OPMX

Chula Vista, CA 91910

PHONE: 619-600-5632

Made for:

inov FARMACEUTIQUE

PRINCIPAL DISPLAY PANEL - 24 Caplets

NDC 69729-601-80

DOLODOL

Alivio temporal de molestias y dolores menores de:

Fiebre, Cabeza y Dental

Fever and temporarily relieves minor aches and pain due to:

Headache-Toothache-Minor pain

24 Caplets

Fever and temporarily relieves
minor aches and pain due to:
Headache-Toothache-Minor pain

DOLODOL

NDC 69729-601-80

DOLODOL

Fever and temporarily relieves
minor aches and pain due to:
Headache-Toothache-Minor pain



Alivio temporal de molestias
y dolores menores de:
Fiebre, Cabeza y Dental



Exclusively distributed by:
OPMX
Chula Vista, CA 91910
Phone: 619-600-5632



Made for:
inov
FARMACEUTIQUE



Drug Facts

Activo Ingrediente (in each caplet)	Purpose
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)	Pain reliever
Caffeine 65 mg	Pain reliever aid
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - cold or arthritis
 - muscular aches
 - toothache
 - menstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes its behavior with nausea and vomiting, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives (red, itchy skin)
- swelling of the face
- difficulty breathing

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning, anticoagulant or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take 3 or more alcoholic drinks every day while using this product (take more or a longer time than directed)

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use if: you ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if:

- you are taking a prescription drug for diabetes, gout, or arthritis
- you are taking any other drug or are under a doctor's care for any serious condition
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

- pregnant or breastfeeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless explicitly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- Keep out of reach of children.
- In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults: 1 full glass of water with each dose
- adults and children 12 years and over:
 - take 2 tablets every 6 hours
 - do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- Store at 20°C (77°F); excursions permitted between 15°C (59°F) and 25°C (77°F).
- TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL, INDICATOR CAP IS BROKEN OR MISSING**

Inactive ingredients

cellulose, croscarmellose, hydroxypropyl methylcellulose, polyvinylpyrrolidone, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

DOLODOL

Fever and temporarily relieves
minor aches and pain due to:
Headache-Toothache-Minor pain

NDC 69729-601-80

DOLODOL

Fever and temporarily relieves
minor aches and pain due to:
Headache-Toothache-Minor pain



Alivio temporal de molestias
y dolores menores de:
Fiebre, Cabeza y Dental



20 Sachets of 4 Caplets

TAMPER EVIDENT:
Do not use if package is
opened or if blister unit is
torn, broken or shows
any sign of tampering

NO VARNISH

Drug Facts

Ingrediente activo (en cada capleta)	Propósito
Acetaminofén 250 mg	Alivio del dolor
Aspirina 250 mg (AINE)	Alivio del dolor
Cafeína 65 mg	Ayuda para el alivio del dolor
*Medicamento antiinflamatorio no esteroideo	

Usos

- Alivia temporalmente dolores y molestias leves debidos a:
 - Dolor de cabeza
 - Rheumatismo
 - Artritis
 - Dolores musculares
 - Síndrome
 - Dolor de muelas
 - Dolores menstruales y menstruales

Advertencias

Síndrome de Reye: Los niños y adolescentes que tienen o se están recuperando de varicela o síntomas similares a la gripe no deben usar este producto. Si al usar este producto hay cambios de comportamiento acompañados de náuseas y vómitos, consulte a un médico, ya que estos síntomas podrían ser un signo temprano del síndrome de Reye, una enfermedad rara pero grave.

Alerta de alergia: La aspirina puede causar una reacción alérgica grave que puede incluir:

- urticaria (erupción)
- hinchazón facial
- Astma (Sibilancias)

Advertencia hepática: Este producto contiene acetaminofén. Puede causar daño hepático grave si usted:

- Toma más de 8 capuletas en 24 horas, que es la cantidad máxima diaria
- Toma con otros medicamentos que contienen acetaminofén
- Consumen 3 o más bebidas alcohólicas al día mientras usa este producto

Advertencia de sangrado estomacal: Este producto contiene un AINE, lo que puede causar sangrado estomacal grave. El riesgo es mayor si usted:

- Tiene 60 años o más
- Ha tenido úlceras estomacales o problemas de sangrado
- Toma un anticoagulante o un esteroide
- Toma otros medicamentos que contienen AINEs (aspirina, ibuprofeno, naproxeno, etc.)
- Consumen 3 o más bebidas alcohólicas al día mientras usa este producto
- Toma más o por más tiempo del indicado

Advertencia sobre la cafeína: La dosis recomendada contiene tanta cafeína como una taza de café. Limite el uso de medicamentos, alimentos o bebidas que contengan cafeína mientras usa este producto, ya que la cafeína puede causar nerviosismo, insomnio, irritabilidad, y ocasionalmente, ritmo cardíaco acelerado.

No use si:

- Es alérgico al acetaminofén, aspirina o a cualquier otro medicamento
- Con cualquier otro medicamento que contenga acetaminofén (recetado o no). Si no está seguro si un medicamento contiene acetaminofén, consulte a un médico o farmacéutico.

Consulte a un médico antes de usar si:

- Tiene enfermedad hepática o la presencia de sangrado estomacal se aplica a usted
- Tiene antecedentes de problemas estomacales, como úlceras
- Tiene presión arterial alta, enfermedad cardíaca, cirrosis hepática o enfermedad renal
- Tiene asma
- Está tomando un diurético

Consulte a un médico o farmacéutico antes de usar si está:

- Tomando un medicamento recetado para diabetes, guta o artritis
- Tomando cualquier otro medicamento o herból o estudio de un médico por una afección seria

Suspenda el uso o consulte a un médico si:

- Experimenta alguno de los siguientes signos de sangrado estomacal:
 - Se siente débil o heces negras o con sangre
 - Vómitos con sangre
 - Dolor estomacal persistente
 - Ocurre una reacción alérgica
- Aspirina muestra síntomas de toxicidad (dolor o ardor de estómago)
- El dolor empeora o dura más de 10 días
- La fiebre empeora o dura más de 3 días
- El área dolorida está roja e hinchada

Si está embarazada o en período de lactancia, consulte a un profesional de salud antes de usar. Es especialmente importante no usar aspirina durante los últimos 3 meses de embarazo, a menos que le indique un médico, ya que puede causar problemas al bebé al nacer o complicaciones durante el parto.

Mantener fuera del alcance de los niños.

En caso de sobredosis, obtenga ayuda médica o contacte al Centro de Control de Envenenamiento (1-800-222-1222) de inmediato. La atención médica rápida es esencial tanto para adultos como para niños, incluso si no se notan signos o síntomas.

Instrucciones

- No usar más de lo indicado
- Beber un vaso lleno de agua con cada dosis
- Adultos y niños de 12 años o más:
 - Tome 2 capuletas cada 6 horas
 - No tome más de 8 capuletas en 24 horas
 - Niños menores de 12 años: consulte a un médico

Otra información


- Almacene a 25°C (77°F); se permite excursion entre 15°C (59°F) and 25°C (77°F).
- EVITAR EL USO DE LA TAPA ESTÁ ROTADO D AUSENTE**
- SEGURIDAD IMPRESO DEBAJO DE LA TAPA ESTÁ ROTADO D AUSENTE**

Ingredientes inactivos:

Artritis de muel, croscarmellose, argemone, celulosa microcristalina, polidona, propilenglicol, lauril sulfato de sodio, glicolato de amoníaco sódico, ácido esteárico, óxido de titanio

NO VARNISH

Pain Reliever / Analgésico
4 Caplets



DOLODOL

DOLODOL

Drug Facts

Active ingredient (in each caplet)

Purpose

Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

* nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
- headache ■ a cold ■ arthritis ■ muscular aches
- sinusitis ■ toothache ■ premenstrual and menstrual cramps

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over
 - take 2 tablets every 6 hours
 - do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

DOLODOL

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-601
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Oblong shape)	Size	17mm
Flavor		Imprint Code	S53

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-601-80	20 in 1 CARTON	01/30/2026	
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/30/2026	

Labeler - OPMX LLC (029918743)**Establishment**

Name	Address	ID/FEI	Business Operations
HBC HEALTHCARE PRIVATE LIMITED		854255181	manufacture(69729-601)

Revised: 1/2026

OPMX LLC