

ACETAMINOPHEN- acetaminophen liquid
Oncor Pharmaceuticals

Acetaminophen

Drug Facts

Active ingredient (in each 5 mL teaspoonful)

Acetaminophen, USP 160 mg

Purpose

Pain reliever/Fever reducer

Uses

For the temporary relief of minor aches and pains associated with:

- the common cold
- flu
- sore throat
- headache
- toothache
- and to reduce fever

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning

drug warfarin.

Stop use and ask a doctor if• pain gets worse or lasts more than 5 days

- pain gets worse or lasts more than 5 day.
- fever gets worse or lasts more than 3 days
- redness or swelling is present.
- any new symptom appear.

These could be signs of a serious condition.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- Use only the enclosed dosing cup designed for use with this product.
- if possible, use weight to dose, otherwise use age.
- dose may be repeated every 4 hours, while symptoms persist, up to five times a day or as directed by a doctor.

Weight (lbs.)	Age (years)	Dosage Teaspoonful (tsp)*
Under 24	Under 2	Consult a doctor
24 to 35	2 to under 4	5 mL (1 tsp)
36 to 47	4 to under 6	7.5 mL (1 1/2 tsp)
48 to 59	6 to under 9	10 mL (2 tsp)
60 to 71	9 to under 11	12.5 mL (2 1/2 tsp)
72 to 95	11 to under 12	15 mL (3 tsp)

* or as directed by a doctor

Other information

- Each 5 mL contains: Sodium 3 mg
- If dispensed, dispense in a tight, light resistant container with a child-resistant cap.
- Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F)

Inactive ingredients

Cherry Flavor, Citric Acid, FD&C Red No. 40, Glycerin, Polyethylene Glycol, Purified Water, Sodium Benzoate, Sodium Citrate, Sodium Saccharin, Sorbitol Solution, Sucralose.

Questions?

You may report side effects by calling Oncor Pharmaceuticals (9 a.m. to 5 p.m. EST), at 1-443-876-7600 or FDA at 1-800-FDA-1088.

Tamper-Evident:

Do not use if foil seal over bottle opening is torn, broken, or missing.

Manufactured in USA For :

Oncor Pharmaceuticals
6755 Business Pkwy Suite 202
Elkridge, MD 21075
United States.

Rev. 01/2024

PRINCIPAL DISPLAY PANEL

ONCOR PHARMACEUTICALS

NDC 83720- 500-16

**Pain and
Fever Relief**

kids

**Acetaminophen
Liquid 160 mg / 5 mL**

Alcohol Free

Sugar Free

Aspirin Free

Ibuprofen Free

Cherry Flavor

16 fl. oz. - 473 mL

ONCOR PHARMACEUTICALS

NDC 83720- 500-16

**Pain and
Fever Relief**

kids

**Acetaminophen
Liquid 160 mg / 5 mL**

Alcohol Free

Sugar Free

Aspirin Free

Ibuprofen Free

Cherry Flavor

4 fl. oz. - 118 mL

NDC 83720-500-16

NCOR
PHARMACEUTICALS

Pain and Fever Relief

Kids

Acetaminophen Liquid

160 mg / 5 mL

Alcohol Free

Sugar Free

Aspirin Free

Ibuprofen Free



NET WT. 16 fl. oz. (473 ml)

DRUG FACTS
Active ingredient (in each 5 mL teaspoonful) Purpose
Acetaminophen, USP 160 mg.....Pain reliever/
fever reducer

Uses
For the temporary relief of minor aches and pains associated with

- the common cold
- flu
- sore throat
- headache
- toothache
- and to reduce fever

Warnings
Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if your child has liver disease.
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.
Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear.

 These could be signs of a serious condition.
Tamper-Evident:
Do not use if foil seal over bottle opening is torn, broken or missing.

DRUG FACTS (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- if possible, use weight to dose, otherwise use age.
- dose may be repeated every 4 hours, while symptoms persist, up to five times a day or as directed by a doctor.

Weight (lbs.)	Age (years)	Dosage Teaspoonful (tsp)*
Under 24	Under 2	Consult a doctor
24 to 35	2 to under 4	5 mL (1 tsp)
36 to 47	4 to under 6	7.5 mL (1 1/2 tsp)
48 to 59	6 to under 9	10 mL (2 tsp)
60 to 71	9 to under 11	12.5 mL (2 1/2 tsp)
72 to 95	11 to under 12	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL contains: Sodium 3 mg
- if dispensed, dispense in a tight, light resistant container with a child-resistant cap.
- store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F)

Inactive ingredients
cherry flavor, citric acid, FD&C red No. 40, glycerin, polyethylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose.

Questions?
You may report side effects by calling Onco Pharmaceuticals (9 a.m. to 5 p.m. EST), at 1-443-876-7600 or FDA at 1-800-FDA-1088.

Manufactured in USA For :
Onco Pharmaceuticals
6755 Business Pkwy Suite 202
Elkridge, MD 21075
United States.
Rev. 12/24

Lot: _____
Exp: _____

NDC 83720-500-04

NCOR

Pain and Fever Relief

Kids

Acetaminophen Liquid

160 mg / 5 mL

Alcohol Free

Sugar Free

Aspirin Free

Ibuprofen Free



NET WT. 4 fl. oz. (118 ml)

Drug Facts
Active ingredient (in each 5 mL teaspoonful) Purpose
Acetaminophen, USP 160 mg.....Pain reliever/fever reducer

Other information

- each 5 mL contains: Sodium 3 mg
- if dispensed, dispense in a tight, light resistant container with a child-resistant cap.
- store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F)

Tamper-Evident:
Do not use if foil seal over bottle opening is torn, broken or missing.

Manufactured in USA For:
Onco Pharmaceuticals
6755 Business Pkwy Suite 202
Elkridge, MD 21075
United States
Rev. 12/24

Drug Facts (continued)

Warnings
Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if your child has liver disease.
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.
Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear.

 These could be signs of a serious condition.

Inactive ingredients
cherry flavor, citric acid, FD&C red No. 40, glycerin, polyethylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose.

Uses
For the temporary relief of minor aches and pains associated with:

- the common cold
- flu
- sore throat
- headache
- toothache
- and to reduce fever

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Questions?
You may report side effects by calling Onco Pharmaceuticals (9 a.m. to 5 p.m. EST), at 1-443-876-7600 or FDA at 1-800-FDA-1088.

Directions

- this product does not contain directions or complete warnings for adult use
- if possible, use weight to dose, otherwise use age.
- dose may be repeated every 4 hours, while symptoms persist, up to five times a day or as directed by a doctor.

Weight (lbs.)	Age (years)	Dosage Teaspoonful (tsp)*
Under 24	Under 2	Consult a doctor
24 to 35	2 to under 4	5 mL (1 tsp)
36 to 47	4 to under 6	7.5 mL (1 1/2 tsp)
48 to 59	6 to under 9	10 mL (2 tsp)
60 to 71	9 to under 11	12.5 mL (2 1/2 tsp)
72 to 95	11 to under 12	15 mL (3 tsp)

* or as directed by a doctor

Manufactured in USA For:
Onco Pharmaceuticals
6755 Business Pkwy Suite 202
Elkridge, MD 21075
United States
Rev. 12/24

ACETAMINOPHEN

acetaminophen liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83720-500
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83720-500-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/10/2024	
2	NDC:83720-500-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/10/2024	

Labeler - Oncor Pharmaceuticals (119032580)

Registrant - Oncor Pharmaceuticals (119032580)