

**COCCIAID- amprolium solution**  
**Aurora Pharmaceutical, Inc.**

-----  
**CocciAid®**

**(amprolium oral solution)**

**9.6% Oral Solution**

**Coccidiostat**

**Water-soluble treatment for coccidiosis**

**Treats 800 gallons at 0.012% level**

Approved by FDA under ANADA # 200-630

**Net Contents:**

1 Gallon (128 fl oz) (3.785 L)

**Active Ingredient:**

amprolium.....9.6%

**INDICATIONS FOR USE:**

CocciAid® (amprolium oral solution) 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

**USE DIRECTIONS:**

Give amprolium at the 0.012% level (8 fl oz per 50 gallons) as soon as coccidiosis is diagnosed and continue for 3 to 5 days. (In severe outbreaks, give amprolium at the 0.024% level.) Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time. Use as the sole source of amprolium. To disperse particles in the product, shake the bottle prior to each use.

**USER SAFETY WARNING:**

Keep this and all drugs out of reach of children.  
NOT FOR HUMAN USE.

**PRECAUTIONS:**

**FOR ORAL USE IN ANIMALS ONLY.**

**MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.

**Restricted Drug (California) — Use Only as Directed**

**STORAGE:**

**Store between 5°-25°C (41°-77°F) with brief excursions to 40°C.**

**Benzoic acid 0.1% added as preservative.**

**CONTACT INFORMATION:**

To report suspected adverse drug experiences, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), contact Aurora Pharmaceutical at 1-888-215-1256 or [www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae)

**REORDER NO: 24001**

**Aurora Pharmaceutical, Inc.**

NORTHFIELD, MN 55057

**1-888-215-1256**

**[www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com)**



CocciAid® is a registered trademark of Aurora Pharmaceutical, Inc.

**FOR ANIMAL USE ONLY**

To Prepare 50 Gallons of Medicated Water

---

**DOSAGE LEVEL MIXING DIRECTIONS**

- 0.024% Add 1 pint (16 fluid ounces) of CocciAid® (amprolium oral solution) 9.6% Oral Solution to about 5 gallons of water in a 50-gallon medication barrel. Stir, then add water to the 50-gallon mark. *Stir thoroughly.*
- 0.012% Follow same directions as above but use 1/2 pint (8 fluid ounces) of CocciAid® 9.6% Oral Solution.
- 0.006% Follow same directions as above but use 4 fluid ounces of CocciAid® 9.6% Oral Solution.
- 

**For Automatic Water Proportioners**

For automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

---

<b>DOSAGE LEVEL</b>	<b>COCCIAID® 9.6% ORAL SOLUTION PER GALLON OF STOCK SOLUTION</b>
0.024%	41 fl oz
0.012%	20.5 fl oz
0.006%	10.25 fl oz

---

**Note:** Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

**Container Label**

PRINCIPAL DISPLAY PANEL - 1 gallon bottle label

IN 50-1465 09/2024

**INDICATIONS FOR USE:** CoccidAid® (amprolium oral solution) 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

**USE DIRECTIONS:** Give amprolium at the 0.012% level (8 fl oz per 50 gallons) as soon as coccidiosis is diagnosed and continue for 3 to 5 days. (In severe outbreaks, give amprolium at the 0.024% level.) Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time. Use this as the sole source of amprolium. To disperse particles in the product, shake the bottle prior to each use.

**WITHDRAWAL PERIODS:** No withdrawal period is required when used according to labeling.

**USER SAFETY WARNING:** Keep this and all drugs out of the reach of children. NOT FOR HUMAN USE.

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY.**  
**MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.  
**Restricted Drug (California) - Use Only as Directed**

**STORAGE:** Store between 5°-25°C (41°-77°F) with brief excursions to 40°C. Benzocaine 0.1% added as preservative.

**CONTACT INFORMATION:** To report suspected adverse drug experiences, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), contact Aurora Pharmaceutical at 1-888-215-1256 or www.aurorapharmaceutical.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VEIS or www.fda.gov/reportanimal

REORDER NO: 24001  
 Aurora Pharmaceutical, Inc.  
 NORTHFIELD, MN 55057  
 1-888-215-1256  
 www.aurorapharmaceutical.com  
 IN 50-1465 09/2024



3 51072 09000 1

Lot  
Exp

TAKE TIME  
OBSERVE LABEL DIRECTIONS




NDC 51072-090-00

**CoccidAid®**  
**(amprolium oral solution)**  
**9.6% Oral Solution**  
**Cocciostat**

Water-soluble treatment for coccidiosis  
 Treats 800 gallons at 0.012% level

Approved by FDA under ANADA # 200-630  
**Net Contents:**  
 1 Gallon (128 fl. oz) (3.785 L)  
**Active Ingredient:**  
 amprolium.....9.6%

**FOR ANIMAL USE ONLY**  
 To Prepare 50 Gallons of Medicated Water

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Add 1 pint (16 fluid ounces) of CoccidAid® (amprolium oral solution) 9.6% Oral Solution to about 5 gallons of water in a 50-gallon medication barrel. Stir, then add water to the 50-gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use 1/2 pint (8 fluid ounces) of CoccidAid® 9.6% Oral Solution.
0.006%	Follow same directions as above but use 4 fluid ounces of CoccidAid® 9.6% Oral Solution.

**For Automatic Water Proportioners**  
 For automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

DOSAGE LEVEL	COCCIDAID® 9.6% ORAL SOLUTION PER GALLON OF STOCK SOLUTION
0.024%	41 fl oz
0.012%	20.5 fl oz
0.006%	10.25 fl oz

Note: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

CoccidAid® is a registered trademark of Aurora Pharmaceutical, Inc.

COCCIDAID			
amprolium solution			
Product Information			
<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:51072-090
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	AMPROLIUM (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	AMPROLIUM	96 in 1 mL
Inactive Ingredients			
	<b>Ingredient Name</b>	<b>Strength</b>	
	WATER (UNII: 059QF0KO0R)		
Packaging			
#	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b> <b>Marketing End Date</b>
1	NDC:51072-090-00	3785 mL in 1 BOTTLE, PLASTIC	
2	NDC:51072-090-01	946 mL in 1 BOTTLE, PLASTIC	
Marketing Information			
	<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b> <b>Marketing End Date</b>
	ANADA	ANADA200630	08/23/2018

**Labeler** - Aurora Pharmaceutical, Inc. (832848639)

## Establishment

Name	Address	ID/FEI	Business Operations
Aurora Pharmaceutical, Inc.		832848639	manufacture

Revised: 10/2024

Aurora Pharmaceutical, Inc.