

**ULTRA BISMUTH- bismuth subsalicylate liquid**  
**Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredient (in each 15 mL)**

Bismuth subsalicylate 525 mg

**Purpose**

Upset stomach reliever/Antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - gas
  - indigestion
  - nausea
  - fullness
  - belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are**

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

### **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
  - 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
  - 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

### **Other information**

- **each 15 mL contains:** sodium 6 mg
- **each 15 mL contains:** salicylate 227 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

## Questions or comments?

Call 1-888-309-9030

## Principal Display Panel

ULTRA

Bismuth

Bismuth subsalicylate 525 mg per 15 mL

Upset stomach reliever / antidiarrheal

Soothing relief for:

- Indigestion
- Upset stomach
- Diarrhea
- Heartburn
- Nausea

2x Strength per ounce<sup>†</sup>

Alcohol free • Sugar free

GLUTEN FREE

FL OZ (mL)

**TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

MANUFACTURED FOR DOLGENCORP, LLC

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072 USA

## Package Label

**Drug Facts** (continued)

**Directions**

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter ■ shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over:
  - 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
  - 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

**Other information**

- each 15 mL contains: sodium 6 mg ■ each 15 mL contains: salicylate 227 mg
- low sodium ■ keep tightly closed ■ protect from freezing
- avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

**Questions or comments?** Call toll free 1-866-4-REXALL

Visit us at: [Rexall.com](http://Rexall.com) or call 1-866-4-REXALL

MANUFACTURED FOR DOLGENCORP, LLC  
100 MISSION RIDGE, GOODLETTSVILLE, TN 37072 USA  
Since 1903 Rexall® is a registered trademark of Rexall Sundown, Inc.

vs. Original Pepto-Bismol®

**GLUTEN FREE**

A0211

3 49580 84212 0

PLD-A559A LB006302

Since 1903  
**Rexall**®

**ULTRA**

# Bismuth

Bismuth subsalicylate 525 mg per 15 mL  
Upset stomach reliever / Antidiarrheal

Soothing relief for:

- Indigestion
- Upset stomach
- Diarrhea
- Heartburn
- Nausea

2x Strength per ounce!  
Alcohol free • Sugar free  
**12 FL OZ (355 mL)**

**Drug Facts** TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

**Active ingredient (in each 15 mL)** **Purpose**  
Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal

**Uses** relieves ■ travelers' diarrhea ■ diarrhea  
■ upset stomach due to overindulgence in food and drink, including  
■ heartburn ■ indigestion ■ nausea  
■ gas ■ fullness ■ belching

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert: Contains salicylate. Do not take if you are**  
■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

**Do not use if you have**  
■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have**  
■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**  
■ diabetes ■ gout ■ arthritis  
■ anticoagulation (thinning the blood)

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**  
■ symptoms get worse or last more than 2 days  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## REXALL Ultra Bismuth

# ULTRA BISMUTH

bismuth subsalicylate liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:559 10-843
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
SODIUM SALICYLATE (UNII: WIQ1H85SYP)	
SORBIC ACID (UNII: X045WJ989B)	
METHYLCELLULOSE (1500 CPS) (UNII: P0NTE48364)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-843-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2018	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part335	09/30/2018	

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 12/2019

Dolgencorp, Inc. (DOLLAR GENERAL &amp; REXALL)