

**NIGHT-TIME MULTI-SYMPTOM COLD/FLU RELIEF- acetaminophen,  
dextromethorphan hydrobromide, doxylamine succinate liquid  
Discount Drug Mart**

-----  
**Cherry Night-Time Multi-Symptom Cold/Flu Relief**

***Drug Facts***

***Active ingredients (in each 30 mL dose cup)***

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine Succinate 12.5 mg

**Purpose**

Pain reliver/fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves cold/flu symptoms

- sore throat
- headache
- minor aches and pain
- fever
- runny nose and sneezing
- cough due to minor throat and bronchial irritation

***Warnings***

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ● skin reddening ● blisters ● rash

If skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription) If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are now taking a prescription monamine oxidase inhibitor (MAOI) (certain drugs for depression psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

#### **Ask a doctor before use if you have**

- a sodium restricted diet
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland.

#### **Ask a doctor or pharmacist before use**

- if you are taking sedatives or tranquilizers
- if you are taking the blood thinning drug warfarin.

#### **When using this product**

- **do not use more than directed**
- avoid alcoholic drinks
- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

#### **Stop use and ask a doctor if**

- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

#### **If pregnant or breast-feeding,**

ask a health professional before use.

#### **Keep out of reach of children.**

#### **Overdose warning:**

Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help, or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms

### Directions

- take only as recommended-see Overdose warning
- Use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hours
- if taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours.

adults and children 12 years and over	30 mL (2TBSP) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

### Other information

- **each 30mL dose cup contains:** sodium 45 mg
- store at room temperature

### Inactive ingredients

citric acid, FD and C Blue No. 1, FD and C Red No. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

### Questions?

Call weekdays **1-877-798-5944**

**DISCOUNT drug mart FOOD FAIR Cherry Night-Time Multi-Symptom Cold/Flu Relief product label**

**Drug Facts (continued)**

Stop use and ask a doctor if • redness or swelling is present  
 • symptoms do not get better within 7 days or are accompanied by a fever  
 • fever gets worse or lasts more than 3 days • new symptoms occur  
 • cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • take only as recommended—see Overdose warning  
 • Use dose cup or tablespoon (TBS) • do not exceed 4 doses per 24 hours  
 • If taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours.

adults & children 12 years and over	30 mL (2 TBS) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

**Other information** • each 30 mL dose cup contains: sodium 45 mg  
 • store at room temperature

**Inactive ingredients** citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

**Questions?** Call weekdays 1-877-798-5944  
 \*This product is not manufactured or distributed by Procter & Gamble owner of the registered trademark Vicks® Nyquil®

**SATISFACTION GUARANTEED**  
 IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LR-056 REV02

0 93351 01759 9

**DISCOUNT**  
**drug mart**  
**FOOD FAIR**

\*COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® NYQUIL®

**Cherry**  
**Night-Time**  
**Multi-Symptom Cold/Flu Relief**

Acetaminophen.....Pain Reliever /Fever Reducer  
 Dextromethorphan HBR..... Cough Suppressant  
 Doxylamine Succinate.....Antihistamine

12 FL OZ (354 mL)

**DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN**  
 Failure to follow these warnings could result in serious consequences

**Drug Facts**

**Active ingredients (in each 30 mL dose cup) Purpose**  
 Acetaminophen 650 mg Pain reliever/fever reducer  
 Dextromethorphan HBR 30 mg Cough suppressant  
 Doxylamine Succinate 12.5 mg Antihistamine

**Uses** temporarily relieves cold/flu symptoms: • sore throat • headache  
 • minor aches and pain • fever • runny nose and sneezing  
 • cough due to minor throat and bronchial irritation

**Warnings**  
**Liver Warning:** The product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using the product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash  
 If skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleepy

**Ask a doctor before use if you have** • a sodium restricted diet • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland.

**Ask a doctor or pharmacist before use** • if you are taking sedatives or tranquilizers • if you are taking the blood thinning drug warfarin

**When using this product** • do not use more than directed  
 • avoid alcoholic drinks • excitability may occur, especially in children  
 • marked drowsiness may occur  
 • be careful when driving a motor vehicle or operating machinery  
 • alcohol, sedatives, and tranquilizers may increase drowsiness

\*COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® NYQUIL®

**DISCOUNT**

**drug mart**

**FOOD FAIR**

**Cherry**

**Night-Time Multi-Symptom Cold/Flu Relief**

Acetaminophen ..... Pain Reliever/Fever Reducer

Dextromethorphan HBR ..... Cough Suppressant

Doxylamine Succinate ..... Antihistamine

12 FL OZ (354 mL)

\* This product is not manufactured or distributed by Procter and Gamble owner of the registered trademark Vicks® Nyquil®

**LR-066 REV 01**

**SATISFACTION GUARANTEED**

IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO THE STORE WHERE PURCHASED. IF UNABLE TO RETURN TO THE STORE, SEND REASON FOR DISSATISFACTION , NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE MEDINNA, OHIO 44256

res

**NIGHT-TIME MULTI-SYMPATOM COLD/FLU RELIEF**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-502
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 4</b> (UNII: X3W0AM1JLX)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SUCROSE</b> (UNII: C151H8M554)	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	CHERRY (CHERRY)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-502-28	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/19/2014	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/19/2014	

**Labeler** - Discount Drug Mart (047741335)

Revised: 12/2024

Discount Drug Mart