

**NIGHT-TIME MULTI-SYMPTOM COLD/FLU RELIEF- acetaminophen,
dextromethorphan hydrobromide, doxylamine succinate liquid
Discount Drug Mart**

Cherry Night-Time Multi-Symptom Cold/Flu Relief

Drug Facts

Active ingredients (in each 30 mL dose cup)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine Succinate 12.5 mg

Purpose

Pain reliver/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves cold/flu symptoms

- sore throat
- headache
- minor aches and pain
- fever
- runny nose and sneezing
- cough due to minor throat and bronchial irritation

Warnings

Liver Warning:This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert:Acetaminophen may cause severe skin reactions. Symptoms may include: ● skin reddening ● blisters ● rash

If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:If sore throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription) If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- a sodium restricted diet
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland.

Ask a doctor or pharmacist before use

- if you are taking sedatives or tranquilizers
- if you are taking the blood thinning drug warfarin.

When using this product

- **do not use more than directed**
- avoid alcoholic drinks
- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning:

Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help, or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms

Directions

- take only as recommended-see Overdose warning
- Use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hours
- if taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours.

adults and children 12 years and over	30 mL (2TBSP) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

Other information

- **each 30mL dose cup contains:**sodium 45 mg
- store at room temperature

Inactive ingredients

citric acid, FD and C Blue No. 1, FD and C Red No. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

Questions?

Call weekdays **1-877-798-5944**

DISCOUNT drug mart FOOD FAIR Cherry Night-Time Multi-Symptom Cold/Flu Relief product label

Drug Facts (continued)

Stop use and ask a doctor if • redness or swelling is present
 • symptoms do not get better within 7 days or are accompanied by a fever
 • fever gets worse or lasts more than 3 days • new symptoms occur
 • cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as recommended-see Overdose warning
 • Use dose cup or tablespoon (TBS) • do not exceed 4 doses per 24 hours
 • If taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours.

adults & children 12 years and over	30 mL (2 TBS) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

Other information • each 30 mL dose cup contains: sodium 45 mg
 • store at room temperature

Inactive ingredients citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

Questions? Call weekdays 1-877-798-5944
 *This product is not manufactured or distributed by Proctor & Gamble owner of the registered trademark Vicks® NyQuil®

SATISFACTION GUARANTEED
 IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO THE STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LR-066 REV02
 0 93351 01759 9



DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN
 Failure to follow these warnings could result in serious consequences

Drug Facts

Active ingredients (in each 30 mL dose cup) Purpose
 Acetaminophen 650 mg Pain reliever/fever reducer
 Dextromethorphan HBR 30 mg Cough suppressant
 Doxylamine Succinate 12.5 mg Antihistamine

Uses temporarily relieves cold/flu symptoms: • sore throat • headache
 • minor aches and pain • fever • runny nose and sneezing
 • cough due to minor throat and bronchial irritation

Warnings
Liver Warning: The product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product. With other drugs containing acetaminophen 3 or more alcoholic drinks every day while using the product.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash
 If skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleepy

Ask a doctor before use if you have • a sodium restricted diet
 • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
 • trouble urinating due to enlarged prostate gland.

Ask a doctor or pharmacist before use • if you are taking sedatives or tranquilizers • if you are taking the blood thinning drug warfarin

When using this product • do not use more than directed
 • avoid alcoholic drinks • excitability may occur, especially in children
 • drowsiness may occur
 • be careful when driving a motor vehicle or operating machinery
 • alcohol, sedatives, and tranquilizers may increase drowsiness

*COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® NYQUIL®

DISCOUNT

drug mart

FOOD FAIR

Cherry

Night-Time Multi-Symptom Cold/Flu Relief

Acetaminophen Pain Reliever/Fever Reducer

Dextromethorphan HBR Cough Suppressant

Doxylamine Succinate Antihistamine

12 FL OZ (354 mL)

* This product is not manufactured or distributed by Proctor and Gamble owner of the registered trademark Vicks® NyQuil®

LR-066 REV 01

SATISFACTION GUARANTEED

IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO THE STORE WHERE PURCHASED. IF UNABLE TO RETURN TO THE STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE MEDINNA, OHIO 44256

res

NIGHT-TIME MULTI-SYMPATOM COLD/FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-502
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 4 (UNII: X3W0AM1JLX)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY (CHERRY)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-502-28	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/19/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/19/2014	

