

**CHILDRENS IBUPROFEN ORAL SUSPENSION- ibuprofen suspension
DEMOULAS SUPERMARKETS INC.**

Children's Ibuprofen Oral Suspension

Drug Facts

Active ingredient(in each 5 mL)

Ibuprofen 100 mg (NSAID)

Purpose

Pain reliever/fever reducer

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: .

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or less or a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- If the child has ever had an allergic reaction to Ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- do not give more than directed
- shake well before using
- mL= milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

- **each 5 mL contains:**sodium 2 mg
- store between 20-25° C (68-77° F)
- **do not use if carton is opened or printed bottle neckband is broken or missing**
- see bottom panel for lot number and expiration date

Inactive ingredients

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Yellow #10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments?

Call 1-877-798-5944 from 9:30 AM to 4:30 PM EST, Monday-Friday.

Package Labeling

MARKET BASKET ®

“MORE FOR YOUR DOLLAR”

Children's

Ibuprofen

Relieves: • Pain • Fever Reducer

MARKET BASKET ®

“MORE FOR YOUR DOLLAR”

***COMPARE TO**

the active ingredient in

CHILDREN'S MOTRIN ®

SEE NEW WARNINGS

Children's

Ibuprofen

Oral Suspension (NSAID)

100mg per 5mL

Relieves: • Pain

• Fever Reducer

Lasts up to 8 hours

ALCOHOL FREE

Original Berry Flafor

4 FL OZ (118 mL)

Important: Read all product information before using.

Keep this box for important information.

This product is intended for use in children.

****This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin[®].**

DISTRIBUTED BY:

DEMOULAS SUPERMARKETS INC.


875 EAST STREET

TEWKSBURY, MA 01876

BX-090

0 49705 74019 4

Box Label



Drug Facts (continued)

- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition or taking any other drug.

When using this product, take with food or milk. If stomach upset occurs.

- Stop use and ask a doctor if:
- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - has bloody or black stools
 - has stomach pain that does not get better
 - child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - the child does not get any relief within first day (24 hours) of treatment
 - fever or pain gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- ml, = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-85	11 years	15 mL

*or as directed by a doctor

Drug Facts (continued)

- Other information**
- each 5 mL contains: sodium 2 mg
 - store between 20-25°C (68-77°F)
 - do not use if cap is opened or printed bottle neckband is broken or missing
 - see bottom panel for lot number and expiration date

Inactive ingredients

Acetylsalicylic acid, anhydrous citric acid, carbonylmethylcellulose sodium, D&C Yellow #10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polyethylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments?

Call 1-877-798-5844 from 9:30 AM to 4:30 PM EST, Monday-Friday.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

DISTRIBUTED BY:
DEMOULAS SUPERMARKETS INC.
876 EAST STREET
TEWKSBURY, MA 01876



Children's Ibuprofen

Relieves: • Pain • Fever Reducer

ALCOHOL FREE

Original Berry Flavor

4 FL OZ (118 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever
fever reducer

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 5 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke

COATING FREE AREA



COATING FREE AREA

Bottle Label

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)*..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: has had stomach ulcers or bleeding

Purpose



problems takes a blood thinning (anticoagulant) or steroid drug takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) takes more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than

2 days or administer to children under 3 years of age unless directed by doctor. **Do not use** if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer right before or after heart surgery **Ask a doctor before use if** stomach bleeding warning applies to your child child has a history of stomach problems, such as heartburn child has problems or serious side effects from taking pain relievers or fever reducers child has not been drinking fluids child has lost a lot of fluid due to vomiting or diarrhea child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke child has asthma child is taking a diuretic **Ask a doctor or pharmacist before use if the child is** under a doctor's care for any serious condition taking any other drug **When using this product** take with food or milk if stomach upset occurs **Stop use and ask a doctor if** child experiences any of the following signs of stomach bleeding: feels faint vomits blood has bloody or black stools has stomach pain that does not get better child has symptoms of heart problems or stroke: chest pain



LR-133



res

CHILDRENS IBUPROFEN ORAL SUSPENSION

ibuprofen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53942-119
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	BERRY (Original)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53942-119-24	1 in 1 CARTON	08/16/2019	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210602	08/16/2019	

Labeler - DEMOULAS SUPERMARKETS INC. (007869647)

Registrant - AptaPharma Inc. (790523323)

Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(53942-119)

Revised: 12/2023

DEMOULAS SUPERMARKETS INC.