CONTROLLED IODINE- iodine solution Durvet Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Controlled Iodine Spray

Antiseptic/Disinfectant

For Skin Disinfection, Superficial Wounds, Navel Cords

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

INDICATIONS: In cattle, horses, swine and sheep as a topical antiseptic for use prior to surgical procedures such as castraing and docking; for ringworms and foot rot; for application to the naveal of newborn animals; and for minor cuts, teat sores, bruises and abraissions.

DIRECTIONS:

Remove spray cap from 16 oz. container and refill from gallon container. Replace spray cap tightly. Hold container approximately 4 to 6 inches from the area to be sprayed. Point valve at the area to be sprayed. Pull trigger of valve and spray area once lightly. May be repeated daily, if necessary, until the area is healed.

CAUTION:

If redness, irritation or swelling persists or increases, discontinue use and consult a veterinarian. Not for use on burns or in body cavities or deep wounds.

NOTE: When used on or near the teats or udders of dairy animals, the teats and udders should be thoroughly washed before the next milking to prevent contamination of the milk.

INGREDIENTS:

Active Ingredients:

Polyethoxylated nonylpenol iodine complex (provides 1.00% minimum titratable iodine).....5.00%

Inactive Ingredients:

Purified Water and Nonionic Surfactant......95.00%

CAUTION: Harmful or fatal if swallowed. Do not apply to the eyes, mucous membraines or large areas of abraided skin. Avoid inhalation of mist.

STORAGE: Store at controlled room temperature between 10°-30°C (50°-86°F) Keep tightly closed, protect from light.

NDC 30798-466-31

FOR ANIMAL USE ONLY

INDICATIONS: In cattle, horses, swine and sheep as a topical antiseptic for use prior to surgical procedures such as castrating and docking; for ringworm and foot rot; for application to the navel of newborn animals; and for minor cuts, teat sores, bruises and abrasions.

DIRECTIONS: Remove spray cap from 16 oz. container and refill from gallon container. Replace spray cap tightly. Hold container approximately 4 to 6 inches from the area to be sprayed. Point valve at the area to be sprayed. Pull trigger of valve and spray area once lightly. May be repeated daily, if necessary, until the area is healed.

CAUTION: If redness, irritation or swelling persists or increases, discontinue use and consult a veterinarian, Not for use on burns or in body cavities or deep wounds.

NOTE: When used on or near the teats or udders of dairy animals, the teats and udders should be thoroughly washed before the next milking to prevent contamination of milk.

Manufactured for DURYET, INC.

Blue Springs, M0 64014 • www.durvet.com

Controlled lodine Spray

Antiseptic/Disinfectant



KEEP OUT OF REACH OF CHILDREN

INGREDIENTS:

Active Ingredients:

Polyethoxylated nonylphenol iodine complex (provides 1,00% minimum titratable iodine)

Inactive Ingredients:95.00% Purified Water and Nonionic


Surfactant,

CAUTION: Harmful or fatal if swallowed. Do not apply to the eyes, mucous membranes or large areas of abraded skin. Avoid inhalation of mist.

STORAGE: Store at controlled room temperature between 10'-30°C (50'-86°F). Keep tightly closed, protect from light.

Lot No.

Exp. Date



FOR ANIMAL USE ONLY

ISS230/B07

INDICATIONS: In cattle, horses, swine and sheep as a topical antiseptic for use prior to surgical procedures such as castrating and docking; for ringworm and foot rot; for application to the navel of newborn animals; and for minor cuts, teat sores, bruises and abrasions.

DIRECTIONS: Remove spray cap from 16 oz. container and refill from gallon container. Replace spray cap tightly. Hold container approximately 4 to 6 inches from the area to be sprayed, Point valve at the area to be sprayed, Pull trigger of valve and spray area once lightly. May be repeated daily, if necessary, until the area is healed.

CAUTION: If redness, irritation or swelling persists or increases, discontinue use and consult a veterinarian, Not for use on burns or in body cavities or deep wounds.

NOTE: When used on or near the teats or udders of dairy animals, the teats and udders should be thoroughly washed before the next milking to prevent contamination of milk.

Manufactured for: **DURVET, INC.**,

Blue Springs, MO 64014 • www.durvet.com ISS23XB



NDC 30798-466-35

KEEP OUT OF REACH OF CHILDREN

INGREDIENTS:

Active Ingredients:

CAUTION: Harmful or fatal if swallowed. Do not apply to the eyes, mucous membranes or large areas of abraded skin, Avoid inhalation of mist,

STORAGE: Store at controlled room temperature between 10"-30"C (50"-86"F), Keep tightly closed, protect from light.



Lot No.

Exp. Date

CONTROLLED IODINE

iodine solution

Product Information Product Type OTC ANIMAL DRUG Item Code (Source) NDC:30798-466 Route of Administration TOPICAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4) O.01 mg in 1 mL

Inactive Ingredients				
Ingredient Name	Strength			
WATER (UNII: 059QF0KO0R)	0.99 mg in 1 mL			

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:30798-466-31	473 mL in 1 BOTTLE					
2	NDC:30798-466-35	3785 mL in 1 JUG					

Marketing Information						
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
	02/01/2012					
	Application Number or Monograph	Application Number or Monograph Marketing Start Citation Date				

Labeler - Durvet Inc (056387798)

Establishment					
Name	Address	ID/FEI	Business Operations		
Centaur, Inc.		174239087	api manufacture, manufacture, label, pack		

Revised: 10/2024 Durvet Inc