

KOSHER MEDS CHILDRENS PAIN AND FEVER- acetaminophen suspension
Mollec Inc

Active Ingredient

(in each 5 mL = 1 teaspoonful)
Acetaminophen 160 mg

Purpose

Pain Reliever/Fever Reducer

Uses

temporarily ■ reduces fever
■ relieves minor aches and pains due to:
■ the common cold ■ flu ■ headache ■ sore throat
■ toothache

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert

acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **shake well before using**
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1 1/2 tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2 1/2 tsp)
72-95 lbs	11 years	15 mL (3 tsp)

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **Keep carton for full directions for use**
- **each 5 mL (1 tsp) contains:** sodium 3 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, butylparaben, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Product label

NON VARNISH/PRINT AREA

kosher-meds

Children's
Pain & Fever

Acetaminophen
Oral Suspension

THE KOSHER CERTIFIED
CHILDREN'S PAIN & FEVER

DO NOT USE IF PRINTED NECK BAND IS
BROKEN OR MISSING

Drug Facts

Active ingredient *Purpose*
(in each 5 mL - 1 teaspoonful)
Acetaminophen 160 mg... Pain reliever/fever reducer

Uses temporarily ■ reduces fever
■ relieves minor aches and pains due to:
■ the common cold ■ flu ■ headache ■ sore throat
■ toothache

Warnings
Liver warning
This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

Allergy alert
acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning
If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose (see overdose warning)
Stop use and ask a doctor if

■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition.
Keep out of reach of children.

Overdose warning
Taking more than the recommended dose (overdose) ▶

Drug Facts (continued)

may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ this product does not contain directions or complete warnings for adult use.
■ shake well before using
■ mL = milliliter; tsp = teaspoonful
■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
■ if needed, repeat dose every 4 hours while symptoms last
■ do not give more than 5 times in 24 hours
■ do not give for more than 5 days unless directed by a doctor.

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-95	11 years	15 mL (3 tsp)

* or as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information
■ Keep carton for full directions for use
■ each 5 mL (1 tsp) contains: sodium 3 mg
■ store between 20-25°C (68-77°F)
■ do not refrigerate

Inactive ingredients anhydrous citric acid, butylparaben, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.
Distributed by: kosher-meds™ BX-155
Mansey, NY 10952 USA • kosher-meds.com

NDC 82179-002-04

COMPARE TO
CHILDREN'S TYLENOL®
ORAL SUSPENSION*
active ingredient

kosher-meds
WITH KOSHER GLYCERIN

Children's
Pain & Fever

Acetaminophen

160 mg per 5mL Oral Suspension
Pain Reliever / Fever Reducer

- Alcohol Free
- Ibuprofen Free
- Aspirin Free



Cherry Flavor

FOR AGES 2 TO 11 4 fl oz (118 mL)

NDC 82179-002-04

COMPARE TO
CHILDREN'S TYLENOL®
ORAL SUSPENSION*
active ingredient

kosher-meds
WITH KOSHER GLYCERIN

Children's
Pain & Fever

Acetaminophen

160 mg per 5mL Oral Suspension
Pain Reliever / Fever Reducer

- Alcohol Free
- Ibuprofen Free
- Aspirin Free



Cherry Flavor

FOR AGES 2 TO 11 4 fl oz (118 mL)



LOT:

EXP:

kosher-meds
WITH KOSHER GLYCERIN

NDC 82179-002-04

**Children's
Pain & Fever**

Acetaminophen

160 mg per 5 mL Oral Suspension
Pain Reliever / Fever Reducer

- Alcohol Free
- Ibuprofen Free
- Aspirin Free

For Ages 2 To 11

Cherry Flavor 4 fl oz (118 mL)

DO NOT USE IF PRINTED NECK BAND IS BROKEN OR MISSING

Active ingredient (in each 5 mL = 1 teaspoonful) **Purpose**
Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses Temporarily ■ reduces fever ■ relieves minor aches and pains due to:
■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings: **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly. Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist. ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product. **Ask a doctor before use if your child has liver disease. Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product do not exceed recommended dose (see overdose warning) Stop use and ask a doctor if ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. Keep out of reach of children. **Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms**

Directions ■ this product does not contain directions or complete warnings for adult use. ■ shake well before using ■ mL = milliliter; tsp = teaspoonful ■ find right dose on chart below. If possible, use weight to dose: otherwise, use age. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give for more than 5 days unless directed by a doctor.

Weight (lb)	Age (yr)	Dose (mL or tsp) *
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-95	11 years	15 mL (3 tsp)

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device. **Other information** ■ each 5 mL (1 tsp) contains: sodium 3 mg ■ store between 20-25°C (68-77°F) ■ do not refrigerate

Inactive ingredients: anhydrous citric acid, butylparaben, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions? Call 1-877-798-5944

Distributed by:

kosher-meds™
Monsey, NY 10952 USA
www.kosher-meds.com



LR-182 **LOT:**

6 **EXP:**

THE KOSHER CERTIFIED CHILDREN'S PAIN & FEVER

KOSHER MEDS CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82179-002
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82179-002-04	1 in 1 CARTON	01/18/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/18/2023	

Labeler - Mollec Inc (040088196)

Revised: 5/2024

Mollec Inc