

## **UTI INFECTION CONTROL- methenamine, sodium salicylate tablet CADENCE HEALTH, INC.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----  
**Drug Facts**

### **Active Ingredient (in each tablet)**

Methenamine 162 mg  
Sodium Salicylate 162.5 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

### **Purpose**

Antibacterial  
Analgesic (pain reliever)

### **Uses**

Temporarily relieves:

- pain and burning
- frequency and urgency of urination

### **Warnings**

**Reye's syndrome:**Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. If changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Stomach bleeding warning:**This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are 60 or older
- have stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do Not Use:**

if you are allergic to salicylates (including aspirin)

**Ask a doctor before use if you have:**

- frequent, burning urination for the first time
- the stomach bleeding warning applying to you
- history of stomach problems, such as heartburn
- high blood pressure
- heart disease
- liver cirrhosis
- bleeding problems
- diuretic use
- ulcers
- kidney disease
- a sodium restricted diet
- reached age 60 or older

**Ask a doctor or pharmacist before use if you are:**

- taking any other drug containing an NSAID (prescription or non-prescription)
- taking a blood thinning (anticoagulant), steroid, diabetes, gout or arthritis drug

**When using this product:**

- do not take more than the recommended dosage

**Stop use and ask a doctor if:**

- product has been used for 3 days
- ringing in the ears
- you experience any of the following signs of stomach bleeding: feel faint
- vomit blood
- have bloody or black stool
- have stomach pain that gets worse or lasts

**If pregnant or breastfeeding**

ask a health professional before use. It is especially important not to use this product (which contains sodium salicylate) during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

***Directions***

**Adults and children 12 years and over:**

- take 2 tablets with a full glass of water 3 times a day. Drink plenty of fluids.
- **Do not use for more than a 3 day period unless directed by a doctor.**

**Children under 12 years:**

- ask a doctor.

**Other information**

- each tablet contains 24.4 mg of sodium
- store at 59-86°F (15-30°C) in a dry place
- protect from light
- Tamper evident: tablets sealed in blisters. Do not use if blister foil or seal is open or damaged.

**Inactive Ingredients**

benzoic acid, croscarmellose sodium, fd&c red #40, fd&c yellow #6, hypromellose, magnesium stearate, methacrylic acid-ethyl acrylate copolymer, microcrystalline cellulose, silicon dioxide, stearic acid, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**MANUFACTURED FOR**

Cadence OTC, Oakland, CA 94612

**Questions or comments:**

Call 1-877-862-7455 or visit us at [cadenceotc.com](http://cadenceotc.com)

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**Cadence**<sup>OTC</sup>

**UTI  
INFECTION  
CONTROL**

Antibacterial to help  
control a UTI

METHENAMINE AND SODIUM SALICYLATE (NSAID)

2 day supply  
12 tablets

NOT INTENDED TO REPLACE MEDICAL CARE.



**Drug Facts**

**Active ingredients** (in each tablet)  
Methenamine 162 mg  
Sodium Salicylate 162.5 mg (NSAID)  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Antibacterial  
Analgesic (pain reliever)

**Uses** Temporarily relieves: • pain and burning • frequency and urgency of urination

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. If changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Stomach bleeding warning:** This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: • are 60 or older • have stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Do not use:** if you are allergic to salicylates (including aspirin)

**Ask a doctor before use if you have:** • frequent, burning urination for the first time • the stomach bleeding warning applying to you • history of stomach problems, such as heartburn • high blood pressure • heart disease • liver cirrhosis • bleeding problems • diuretic use • ulcers • kidney disease • a sodium restricted diet • reached age 60 or older

**Ask a doctor or pharmacist before use if you are:** • taking any other drug containing an NSAID (prescription or non-prescription) • taking a blood thinning (anticoagulant), steroid, diabetes, gout or arthritis drug

**When using this product:** • do not take more than the recommended dosage

**Stop and ask a doctor if:** • product has been used for 3 days • ringing in the ears • you experience any of the following signs of stomach bleeding: feel faint • vomit blood • have bloody or black stool • have stomach pain that gets worse or lasts

**If pregnant or breast feeding,** ask a health professional before use. It is especially important not to use this product (which contains sodium salicylate) during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

**Directions** Adults and children 12 years and over: • take 2 tablets with a full glass of water 3 times a day. Drink plenty of fluids. • Do not use for more than a 3 day period unless directed by a doctor.

**Children under 12 years:** • ask a doctor.

**Other information** • each tablet contains 24.4 mg of sodium • store at 59-96°F (15-30°C) in a dry place • protect from light • Tamper evident: tablets sealed in blisters. Do not use if blister foil or seal is open or damaged.

**Inactive ingredients** benzoic acid, croscarmellose sodium, fd&c red #40, fd&c yellow #6, hypromellose, magnesium stearate, methacrylic acid-ethyl acrylate copolymer, microcrystalline cellulose, silicon dioxide, stearic acid, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**MANUFACTURED FOR** Cadence OTC, Oakland, CA 94612  
**Questions or comments:** Call 1-877-862-7455 or visit us at cadenceotc.com  
**CSDUALCAD** Rev 1/26

**UTI INFECTION CONTROL**

methenamine, sodium salicylate tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73436-1221
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>METHENAMINE</b> (UNII: J50OIX95QV) (METHENAMINE - UNII:J50OIX95QV)	METHENAMINE	162 mg
<b>SODIUM SALICYLATE</b> (UNII: WQ1H85SYP) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SODIUM SALICYLATE	162.5 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKNOB0MIM)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	PH061
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73436-1221-2	1 in 1 CARTON	04/15/2026	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/06/2026	

**Labeler** - CADENCE HEALTH, INC. (116830263)

Revised: 4/2026

CADENCE HEALTH, INC.