

FENBENDAZOLE ORAL DEWORMER- fenbendazole paste
Shenzhen Zhenshangzekai Technology Co., Ltd.

Febendazole

Glycerin, Magnesium Aluminum Silicate, Sodium Carboxymethylcellulose, Sorbitol, and Purified

Water.

Keep out of the reach of children.

The antiparasitic efficacy is achieved through the systematic inhibition of energy metabolism

within the target parasites.

Standard Therapeutic Dose:

2.3 mg/lb(5 mg/kg) of body weight.

Intensive Care Dose:

4.6 mg/lb(10 mg/kg) for specific larval stages or young animals.

Calibration Instructions:

1. Accurately assess the animal's weight prior to administration.
2. Remove the protective tip and rotate the dial ring to the zero position.
3. Advance the dial ring to the graduation corresponding to the measured weight.
4. Ensure the oral cavity is clear of food particles.
5. Administer the paste onto the back of the tongue by depressing the plunger firmly.

Yield: One 25g syringe provides a complete dose for a 1,100 lb animal at the standard 5 mg/kg rate.

dewormer

Human Safety: NOT FOR USE IN HUMANS.

- Food Safety: Do not use in animals intended for human consumption.
- Animal Care: Safety in pregnant animals or breeding stallions at the 10 mg/kg dosage for 5 days has not been specifically evaluated.
- Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Resistance Management:

Fecal examinations and diagnostic monitoring are recommended to maintain long-term efficacy and manage parasite resistance.

Environmental & Storage Conditions:




- Store at or below 25°C (77°F) in a temperature-controlled environment.
- Keep the syringe tightly capped when not in use.

Indicated for the clinical management of:

Large Strongyles: (Strongylus edentatus, S. equinus, S. vulgaris).

Small Strongyles: Including various larval stages (encysted early 3rd stage, late 3rd stage, and 4th stage cyathostome larvae).

Ascarids / Roundworms: (*Parascaris equorum*).
 Pinworms: (*Oxyuris equi*).

<p>SAFETY & REGULATORY</p> <p>Safety Precautions:</p> <ul style="list-style-type: none"> Human Safety: NOT FOR USE IN HUMANS. Food Safety: Do not use in animals intended for human consumption. Animal Care: Safety in pregnant animals or breeding stallions at the 10 mg/kg dosage for 5 days has not been specifically evaluated. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. <p>Resistance Management:</p> <p>Fecal examinations and diagnostic monitoring are recommended to maintain long-term efficacy and manage parasite resistance.</p> <p>Environmental & Storage Conditions:</p> <ul style="list-style-type: none"> Store at or below 25°C (77°F) in a temperature-controlled environment. Keep the syringe tightly capped when not in use. <p>Questions or Comments? Email: PREPAKU2026@outlook.com Call toll-free: 1-877-555-0193 Distributed by: PREPAKU INC. 789 Innovation Way, Snow Hill, NC 28582 MADE IN USA LOT: P-115 EXP: 12/2029</p>	<p>DOSAGE & ADMINISTRATION</p> <p>Standard Therapeutic Dose: 2.3 mg/lb(5 mg/kg) of body weight.</p> <p>Intensive Care Dose: 4.6 mg/lb(10 mg/kg) for specific larval stages or young animals.</p> <p>Calibration Instructions:</p> <ol style="list-style-type: none"> 1. Accurately assess the animal's weight prior to administration. 2. Remove the protective tip and rotate the dial ring to the zero position. 3. Advance the dial ring to the graduation corresponding to the measured weight. 4. Ensure the oral cavity is clear of food particles. 5. Administer the paste onto the back of the tongue by depressing the plunger firmly. <p>Yield: One 25g syringe provides a complete dose for a 1,100 lb animal at the standard 5 mg/kg rate.</p> <p>NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN.</p> 	<p>PREPAKU®</p> <p>FENBENDAZOLE</p> <p>Oral Dewormer</p> <p>25 gram Paste 10% (100 mg/g)</p> <ul style="list-style-type: none"> ✓ Large Strongyles ✓ Small Strongyles ✓ Roundworms ✓ Pinworms  	<p>PRODUCT DESCRIPTION</p> <p>Active Ingredient: Each gram contains 100 mg of Fenbendazole.</p> <p>Inactive Ingredients: Glycerin, Magnesium Aluminum Silicate, Carboxymethylcellulose Sodium, Sorbitol, and Purified Water.</p> <p>Pharmacological Action: The antiparasitic efficacy is achieved through the systematic inhibition of energy metabolism within the target parasites.</p> <p>INDICATIONS FOR USE</p> <p>Indicated for the clinical management of:</p> <p>Large Strongyles: (<i>Strongylus edentatus</i>, <i>S. equinus</i>, <i>S. vulgaris</i>).</p> <p>Small Strongyles: Including various larval stages (<i>encysted early 3rd stage, late 3rd stage, and 4th stage cyathostome larvae</i>).</p> <p>Ascarids / Roundworms: (<i>Parascaris equorum</i>).</p> <p>Pinworms: (<i>Oxyuris equi</i>).</p>
---	--	---	--

FENBENDAZOLE ORAL DEWORMER

fenbendazole paste

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87276-263
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FENBENDAZOLE (UNII: 621BVT9M36) (FENBENDAZOLE - UNII:621BVT9M36)	FENBENDAZOLE	100 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
AQUA (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
GLYCERIN (UNII: PDC6A3C0OX)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87276-263-00	1 g in 1 TUBE; Type 0: Not a Combination Product	03/17/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M024	03/17/2026	

Labeler - Shenzhen Zhenshangzekai Technology Co., Ltd. (518601007)

Revised: 3/2026

Shenzhen Zhenshangzekai Technology Co., Ltd.