

**ALLERGY MULTI-SYMPTOM- acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet, coated**  
**AMERICAN SALES COMPANY**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**1128 - CAR - 21018-1229**

***Drug Facts***

<b><i>Active ingredients (in each caplet)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 325 mg	Pain reliever
Chlorpheniramine maleate 2 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
  - headache
  - sinus congestion and pressure
  - nasal congestion
  - runny nose and sneezing
  - minor aches and pains
- temporarily relieves these additional symptoms of hay fever:
  - itching of the nose or throat
  - itchy, watery eyes
- helps clear nasal passages
- helps decongest sinus openings and passages

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

#### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

#### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

#### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

#### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than directed (see overdose warning)

---

adults and

- take 2 caplets every 4 hours
- swallow whole - do not crush, chew,

adults and children 12 years and over	<ul style="list-style-type: none"> <li>or dissolve</li> <li>do not take more than 12 caplets in 24 hours</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>ask a doctor</li> </ul>

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

**Inactive ingredients**

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, flavor, hypromellose, iron oxide ochre, lactose anhydrous, magnesium stearate, povidone, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide

**PRINCIPAL DISPLAY PANEL**

NDC 41520-601-02

CareOne®

For Adults

Allergy Multi-Symptom

Pain Reliever – Acetaminophen

Nasal Decongestant - Phenylephrine HCl

Antihistamine – Chlorpheniramine Maleate

May Cause Drowsiness

Instant Cooling Sensation

For the Relief of:

Headache


Sinus Pressure; Nasal Congestion

Runny Nose; Sneezing

Itchy, Watery Eyes

Our Pharmacists Recommend

24 Cool Taste Caplets



CAREone®

For Adults

# ALLERGY MULTI-SYMPTOM

**Pain Reliever - Acetaminophen**  
**Nasal Decongestant - Phenylephrine HCl**  
**Antihistamine - Chlorpheniramine Maleate**

**May Cause Drowsiness**  
**Instant Cooling Sensation**  
**For the Relief of:**  
**Headache**  
**Sinus Pressure, Nasal Congestion**

Actual Size

2017-109-41520-NC

NC

NC

### Drug Facts (continued)

**Active ingredients (in each caplet)**  
 Acetaminophen 325 mg.....Pain reliever  
 Chlorpheniramine maleate 2 mg.....Antihistamine  
 Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses** ■ temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ headache ■ sinus congestion and pressure ■ nasal congestion ■ runny nose and sneezing ■ minor aches and pains ■ helps clear nasal passages ■ helps decongest sinus openings and passages

### Drug Facts (continued)

**Ask a doctor or pharmacist before use if you are**  
 ■ taking sedatives or tranquilizers  
 ■ taking the blood thinning drug warfarin

**When using this product**  
 ■ do not exceed recommended dosage  
 ■ excitability may occur, especially in children  
 ■ drowsiness may occur ■ avoid alcoholic drinks  
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness  
 ■ be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**  
 ■ nervousness, dizziness, or sleeplessness occur  
 ■ pain or nasal congestion gets worse or lasts more than 7 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ new symptoms occur  
 ■ redness or swelling is present

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 ■ do not take more than directed (see overdose warning)  
 adults and children 12 years and over  
 ■ take 2 caplets every 4 hours, swallow whole - do not crush, chew, or dissolve  
 ■ do not take more than 12

### Drug Facts (continued)

■ temporarily relieves these additional symptoms of  
 ■ hay fever: ■ itching of the nose or throat  
 ■ itchy, watery eyes

**Warnings**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 12 caplets in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
 ■ liver disease  
 ■ high blood pressure  
 ■ heart disease  
 ■ diabetes  
 ■ glaucoma



F112802CAR\_R1 S-1237 1117

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

DISTRIBUTED BY FOODHOLD U.S.A., LLC  
 LANDOVER, MD 20785  
 1-877-846-9949  
 © 2017 S&S Brands, LLC  
 Quality guaranteed or your money back.



sinus pressure, nasal congestion  
Runny Nose; Sneezing  
Itchy, Watery Eyes



**24 COOL TASTE CAPLETS**

## ALLERGY MULTI-SYMPTOM

acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet, coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-601
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	white (Off-White)	Score	no score
Shape	OVAL	Size	17mm
Flavor	MINT	Imprint Code	AAA;1117
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-601-02	2 in 1 CARTON	08/01/2011	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/01/2011	

**Labeler** - AMERICAN SALES COMPANY (809183973)

Revised: 12/2018

AMERICAN SALES COMPANY