

**POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution
Cardinal Health 107, LLC**

Drug Facts

Polyethylene Glycol 3350, Powder for Oral Solution

Active Ingredient (in each sachet)

Polyethylene Glycol 3350, NF 17g

Purpose

Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol.

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor of pharmacist

before use if you are taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
- use once a day
- stir and dissolve one sachet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing.
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other Information

- store at 20° - 25°C (68° - 77°F)
- do not use if foil is open or broken.

Inactive Ingredients

none

Questions or comments?

1-800-707-4621 or www.americanhealthpackaging.com

© 2023 & Distributed by
American Health Packaging
Columbus, Ohio 43217 USA

All rights reserved.

Overbagged with 7 x 0.6 OZ (17 g) packets in each bag, NDC 55154-8184-5

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. Keep this and all drugs out of the reach of children.

Distributed By:

Cardinal Health

Dublin, OH 43017

L5568654-10124 / L5568654-20124

1047662/1047664/1049041/1046550

Principal Display Panel

NDC 55154-8184-5

HEALTHYLAX®

POLYETHYLENE GLYCOL 3350 NF

Powder for Oral Laxative Solution

7 SACHETS 0.6 OZ (17g) EACH



NDC 55154-8184-5

HEALTHYLAX®

POLYETHYLENE GLYCOL 3350 NF

Powder For Oral Laxative Solution

7 SACHETS 0.6 OZ (17g) EACH

- Relieves Occasional Constipation/Irregularity
- Softens Stool
- 7 Daily Doses

Drug Facts

Active Ingredient (in each sachet)
Polyethylene Glycol 3350 NF, 17g

Inactive Ingredients: none

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

STORAGE: Store at 20° - 25° C (68° - 77° F)
do not use if foil is open or broken.

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. If pregnant or breast-feeding, ask a health professional before use. Keep this and all medications out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Code No.: KR/DRUGS/KTK/25/415/98

© 2019 & Distributed by American Health Packaging,

Columbus, Ohio 43217 USA. All rights reserved.

Distributed by Cardinal Health
Dublin, OH 43017

L5568654-10124

Lot: Exp:

F106

Distributed by Cardinal Health
Dublin, OH 43017
L5568654-20124

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting, abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome.

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.

- you get diarrhea

- you need to use a laxative for longer than 1 week

Directions

- Do not take more than directed unless advised by your doctor

- Adults and children 17 years of age and older:

- Stir and dissolve one sachet of powder (17g) in any 4 to 8 ounces of beverage (cold, hot or room temperature)

- Drink the solution

- Use once a day

- Use no more than 7 days

- Children 16 years of age or under: ask a doctor

Questions or comments?

1-800-707-4621 or

www.americanhealthpackaging.com

POLYETHYLENE GLYCOL 3350

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55154-8184(NDC:60687-431)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	white (Colorless upon dissolution)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-8184-5	7 in 1 BAG	04/15/2019	
1		17 g in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203928	04/15/2019	

Labeler - Cardinal Health 107, LLC (118546603)

Revised: 10/2025

Cardinal Health 107, LLC