

**SERYNTH NICOTINE TRANSDERMAL STOP SMOKING AID- nicotine21mg
nicotine transdermal stop smoking aid patch
Jiangxi Yudexi Pharmaceutical Co., LTD**

85248-164

Active Ingredient

Nicotine 21mg

Purpose

stop smoking aid

Use

reduces withdrawal symptoms,including nicotine craving,associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine,This medicine is believed to be safer than smoking.However, the risks to your child from this medicine are not fully known

Do not use

Do not wear more than one patch at a time.Do not cut patch in half or into smaller pieces,Do not leave patch on for more than 24 hours.because it may irritate your skin and loses strength after 24 hours.To avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging)procedures.It is important to complete treatment,if you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

When Using

if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop Use

Stop use and ask a doctor if Feeling uncomfortable or experiencing or other sensitive dizziness symptoms

Ask Doctor

Ask a doctor before use if you have Heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. High blood pressure not controlled with medication. Nicotine can increase your blood pressure.

An allergy to adhesive tape or have skin problems because you are more likely to get rashes. Stomach ulcer or diabetes, History of seizures.

Ask a doctor or pharmacist before use if you are: Using a non-nicotine stop smoking drug.

Taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Keep Out Of Reach Of Children

Keep out of reach of children and pets.

Directions

If you are under 18 years of age, ask a doctor before use. Before using this product, need the enclosed self-help guide for complete directions and other information. Begin using the patch on your quit day. If you smoke more than 10 cigarettes per day, use the following schedule below. The used patch should be removed and a new one applied to a different skin site at the same time each day. If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning. Do not wear more than one patch at a time. Do not cut patch in half or into smaller pieces. Do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours. To avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures. It is important to complete treatment, if you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider. Signs of excessive use may include nausea, vomiting, dizziness, weakness, or rapid heartbeat. If these symptoms occur, remove the patch immediately and contact a doctor or poison control center.

Other information

store at 20-25°C

Inactive ingredients

acrylate adhesive □ cellulose paper □ methacrylic acid .

PRINCIPAL DISPLAY PANEL

SERYNTH

35*75*110MM

SERYNTH

NICOTINE TRANSDERMAL STOP SMOKING AID

24-HOUR CRAVING CONTROL

QUIT SMOKING

- 24-hour craving control
- Reduces withdrawal symptoms
- Helps you quit smoking

21mg Per Patch

21 Patches

NICOTINE TRANSDERMAL STOP SMOKING AID

NICOTINE TRANSDERMAL STOP SMOKING AID

Drug Facts

Active ingredient (in each patch) Purpose
Nicotine 21mg stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have Heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. High blood pressure not controlled with medication. Nicotine can increase your blood pressure.

An allergy to adhesive tape or have skin problems because you are more likely to get rashes. Stomach ulcer or diabetes. History of seizures.

Ask a doctor or pharmacist before use if you are: Using a non-nicotine stop smoking drug. Taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product If you have vivid dreams or other sleep disturbances remove this patch at bedtime.

Stop use and ask a doctor if Feeling uncomfortable or experiencing or other sensitive dizziness symptoms.

Keep out of reach of children and pets.

Directions If you are under 18 years of age, ask a doctor before use.

Before using this product, read the enclosed self-help guide for complete directions and other information.

Begin using the patch on your quit day. If you smoke more than 10 cigarettes per day, use the following schedule below.

The used patch should be removed and a new one applied to a different skin site at the same time each day. If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning. Do not wear more than one patch at a time. Do not cut patch in half or into smaller pieces. Do not leave patch on for more than 24 hours, because it may irritate your skin and loses strength after 24 hours.

To avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures. It is important to complete treatment, if you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Signs of excessive use may include nausea, vomiting, dizziness, weakness, or rapid heartbeat. If these symptoms occur, remove the patch immediately and contact a doctor or poison control center.

Inactive Ingredients: acrylate adhesive, cellulose paper, methacrylic acid.
Other information: store at 20° to 25°C (68° to 77°F)



Distributor: Jiangxi Yudexi Pharmaceutical Co., LTD
Address: No.1998, Jingkai West 5th Road, Chengshu Jingkai District, Zhangshu City, Yichun City, Jiangxi Province 331200, China (CN)

SERYNTH NICOTINE TRANSDERMAL STOP SMOKING AID

nicotine21mg nicotine transdermal stop smoking aid patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85248-164
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	21 mg in 100

Inactive Ingredients

Ingredient Name	Strength
2-HYDROXYETHYL ACRYLATE (UNII: 25GT92NY0C)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
METHACRYLIC ACID (UNII: 1CS02G8656)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85248-164-01	21 in 1 BOX; Type 0: Not a Combination Product	03/19/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	03/19/2026	

Labeler - Jiangxi Yudexi Pharmaceutical Co., LTD (455662836)

Revised: 3/2026

Jiangxi Yudexi Pharmaceutical Co., LTD