FOSTER AND THRIVE ASPIRIN- aspirin tablet, chewable Strategic Sourcing Services LLC

McKesson Aspirin Drug Facts

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- · you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- · you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- · drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- · children under 12 years: consult a doctor

Other information

store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

Questions or comments?

Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

Principal Display Panel

COMPARE TO BAYER® CHEWABLE

ACTIVE INGREDIENT

Foster & Thrive[™]

Aspirin

ORANGE FLAVORED

Pain Reliever (NSAID)


Low Dose Aspirin Regimen | Gluten Free

ACTUAL SIZE

Talk to your doctor or other healthcare provider before using this product for your heart.

SEE NEW WARNING

36 Chewable Tablets - 81 mg



NDC 70677-1134-1

Foster & Thrive™ spirin

Pain Reliever (NSAID)

Low Dose Aspirin Regimen | Gluten Free Talk to your doctor or other healthcare provider before using this product for your heart.

DO NOTUSE IF PRINTED S EAL UNDER CAP IS BROKENOR MISSING

SEE NEW WARNING

COMPARE TO BAYER® CHEWABLE **ACTIVE INGREDIENT****

> ORANGE FLAVORED

ACTUAL SIZE

36 Chewable Tablets - 81 mg

Rev.09/23

Dis tri but ed by: McKesson Corp.,
vi a Strategic Sourcing Services LLC.
Memphins, IT & 3844
Morrey Back Guarantee
www. Astern arthin ve.com



Drug Facts

Active ingredient Purpose (in each chewable tablet) ..Pain reliever Aspirin 81 mg (NSAID*).....Pair *nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox orflu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling
- asthma (wheezing) shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)

KEEP OUTER CARTON FOR COMP WARNINGS AND PRODUCTINFORM

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an all ergic reaction to any other pain reliever/fever reducer

■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach blæding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kid ney disease
- you are taking a diuretic you have asthma

Ask a doctor or pharma cist before use if you are taking a prescription drug for

■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint vomit blood have bloody or black stools have stomach pain thatdoes not get better
- pain gets wors e or lasts more than 10 days
- fever gets worse or lasts more than 3 days redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs These could be signs of a serious condition

Drug Facts (continued)

lfpregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Pois on Control Center right away (1-800-222-1222).

- drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours un less directed by a doctor
- children under 12 years: consulta doctor

Other information

store at 20-25°C (68-77°F)

Inactive ingredients com starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

Questions or comments?

Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

**This product is not manufactured or distributed by Bayer Health Care LLC, distributor of Bayer® Chewable.



46768 KM C1

aspirin tablet, chewable

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:70677-1134

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | | Basis of Strength | Strength |
|-----------------|---|--------------------------|----------|
| | ASPIRIN (LINII: R16CO5Y76E) (ASPIRIN - LINII: R16CO5Y76E) | AS PIRIN | 81 mg |

Inactive Ingredients Ingredient Name Strength STARCH, CORN (UNII: 08232NY3SJ) FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q) SACCHARIN SODIUM (UNII: SB8ZUX40TY)

| Product Characteristics | | | | | |
|-------------------------|--------|--------------|----------|--|--|
| Color | ORANGE | Score | no score | | |
| Shape | ROUND | Size | 8mm | | |
| Flavor | ORANGE | Imprint Code | L467 | | |
| Contains | | | | | |

| Packaging | | | | | |
|------------------------|---|-------------------------|-----------------------|--|--|
| # Item Code | Package Description | Marketing Start Date | Marketing End Date | | |
| 1 NDC:70677- 1134-1 | 1 in 1 CARTON | 11/02/2023 | | | |
| 1 | 36 in 1 BOTTLE; Type 0: Not a Combination Product | | | | |

| Marketing Information | | | | | | |
|-----------------------|---|-------------------------|-----------------------|--|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | | |
| OTC Monograph Drug | M013 | 11/02/2023 | | | | |
| | | | | | | |

Labeler - Strategic Sourcing Services LLC (116956644)