

SMILECARE MOUTH ULCER GEL- menthol gel
Yantai Aien Biotechnology Co., Ltd.

87512-004 SmileCare Mouth Ulcer Gel

Active ingredient

Purpose

Menthol(0.1%)..... Oral pain reliever

Uses

Temporarily relieves occasional minor irritation and pain associated with sore mouth and sore throat.

Keep out of reach of children

Apply a small amount of gel to the affected area of the mouth. Gargle, swish around, or allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.

Warnings

Do not use

In children under 2 years of age unless directed by a doctor

Sore throat warnings

1If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. These may be serious.


Directions

Adults and children 2 years of age and older: Apply a small amount of gel to the affected area of the mouth. Gargle, swish around, or allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor. Children under 12 years of age should be supervised in the use of this product. Children under 2 years of age: Consult a dentist or doctor.


Inactive ingredients

Polihexanide, Riboflavin, Ferrous Sulfate, Sorbitol, Sodium Alginate, Glycerin, Niacinamide, Carbomer Homopolymer, Trolamine, Water

87512-004-01 SmileCare Mouth Ulcer Gel 10g



87512-004-02 SmileCare Mouth Ulcer Gel 15g



Eisen Life

NDC 0000000000



EisenLife

**SMILECARE
MOUTH ULCER
GEL**

SOOTHES
IRRITATED GUMS

DELIVERS
TARGETED PAIN
RELIEF

Eisen Life
**FAST ACTING ORAL
PAIN RELIEF**

- Toothache
- Gum Pain
- Canker Sores
- Mouth Irritation



NET WT 0.53 OZ (15 g)

MADE IN CHINA

Drug Facts

Active ingredient	Purpose
Menthol(0.1%),.....	Oral pain reliever

Uses

- Temporarily relieves occasional minor irritation and pain associated with sore mouth and sore throat.

Warnings

Do not use

- In children under 2 years of age unless directed by a doctor

Sore throat warnings

- If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. These may be serious.

Keep out of reach of children

Directions


- Adults and children 2 years of age and older: Apply a small amount of gel to the affected area of the mouth. Gargle, swish around, or allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.
- Children under 12 years of age should be supervised in the use of this product.
- Children under 2 years of age: Consult a dentist or doctor.

Other Information

- Store at room temperature.
- Protect this product from excessive heat and direct sun.

Inactive ingredients

Polihexanide, Riboflavin, Ferrous Sulfate, Sorbitol, Sodium Alginate, Glycerin, Niacinamide, Carbomer Homopolymer, Triethylamine, Water



Eisen Life

NDC 000000000



EisenLife

**SMILECARE
MOUTH ULCER
GEL**

SOOTHES
IRRITATED GUMS

DELIVERS
TARGETED PAIN
RELIEF

Eisen Life
**FAST ACTING ORAL
PAIN RELIEF**

- Toothache
- Gum Pain
- Canker Sores
- Mouth Irritation

NET WT 0.71 OZ (20 g)

MADE IN CHINA



Drug Facts

Active ingredient	Purpose
Menthol(0.1%),.....	Oral pain reliever

Uses

- Temporarily relieves occasional minor irritation and pain associated with sore mouth and sore throat.

Warnings

Do not use

- In children under 2 years of age unless directed by a doctor

Sore throat warnings

- If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. These may be serious.

Keep out of reach of children

Directions

- Adults and children 2 years of age and older: Apply a small amount of gel to the affected area of the mouth. Gargle, swish around, or allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.
- Children under 12 years of age should be supervised in the use of this product.
- Children under 2 years of age: Consult a dentist or doctor.

Other Information

- Store at room temperature.
- Protect this product from excessive heat and direct sun.

Inactive ingredients

Polihexanide, Riboflavin, Ferrous Sulfate, Sorbitol, Sodium Alginate, Glycerin, Niacinamide, Carbomer Homopolymer, Triethylamine, Water

SMILECARE MOUTH ULCER GEL

menthol gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87512-004
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.1 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
GLYCERIN (UNII: PDC6A3C0OX)	
RIBOFLAVIN (UNII: TLM2976OFR)	
TROLAMINE (UNII: 9O3K93S3TK)	
WATER (UNII: 059QF0KO0R)	
CARBOMER HOMOPOLYMER, UNSPECIFIED TYPE (UNII: 0A5MM307FC)	
NIACINAMIDE (UNII: 25X51I8RD4)	
POLIHESANIDE (UNII: 322U039GMF)	
SORBITOL (UNII: 506T60A25R)	
FERROUS SULFATE (UNII: 39R4TAN1VT)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87512-004-01	10 g in 1 TUBE; Type 0: Not a Combination Product	03/20/2026	
2	NDC:87512-004-02	15 g in 1 TUBE; Type 0: Not a Combination Product	03/20/2026	
3	NDC:87512-004-03	20 g in 1 TUBE; Type 0: Not a Combination Product	03/20/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	03/20/2026	

Labeler - Yantai Aien Biotechnology Co., Ltd. (977916686)