MINOXIDIL FOR WOMEN- minoxidil solution Chain Drug Consortium, LLC

Minoxidil Topical Solution for Women

Drug Facts

Active ingredient

Minoxidil, USP 2% w/v

Purpose

Hair regrowth treatment

Use

to regrow hair on the scalp

Warnings

For external use only

Flammable: Keep away from fire or flame

Do not use if

- your degree of hair loss is different than that shown on the side of this carton, because this product may not work for you
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- your hair loss is associated with childbirth
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

Ask a doctor before use if you have

heart disease

When using this product

- do not apply on other parts of the body
- avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. You may need to use this product 2 times a day for at least 4 months before you see results.

• the amount of hair regrowth is different for each person. This product will not work for everyone.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply one mL with dropper 2 times a day directly onto the scalp in the hair loss area
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- in clinical studies of mostly white women aged 18 to 45 years with mild to moderate degrees of hair loss, the following response to 2% minoxidil topical solution was reported: 19% of women reported moderate hair regrowth after using 2% minoxidil topical solution for 8 months (19% had moderate regrowth; 40% had minimal regrowth). This compares with 7% of women reporting moderate hair regrowth after using the placebo, the liquid without minoxidil in it, for 8 months (7% had moderate regrowth; 33% had minimal regrowth).
- store at controlled room temperature 20° to 25°C (68° to 77°F).

Inactive ingredients

alcohol (60% v/v), propylene glycol, purified water

Questions?

1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

Principal Display Panel

Premier Value®

COMPARE TO THE ACTIVE INGREDIENT IN WOMEN'S ROGAINE®*

For Women Minoxidil TOPICAL SOLUTION, USP 2% HAIR REGROWTH TREATMENT

Reactivates Hair Follicles to stimulate regrowth

Clinically Proven to Help Regrow Hair

Unscented

FOR BEST RESULTS
APPLY DIRECTLY TO SCALP
TWICE DAILY

INDEPENDENTLY TESTED SATISFACTION GUARANTEED

Two Month Supply
Two 60 mL (2 fl oz) Bottles





COMPARE TO THE ACTIVE INGREDIENT IN WOMEN'S ROGAINE®*

For Women Minoxidil

TOPICAL SOLUTION, USP 2% HAIR REGROWTH TREATMENT

Reactivates Hair Follicles to Stimulate Regrowth

Clinically Proven to Help Regrow Hair


Unscented

FOR BEST RESULTS

APPLY DIRECTLY TO SCALP
TWICE DAILY

Two Month Supply
Two 60mL (2 fl oz) Bottles









Drug Facts

Active ingredient

Purpose

Minoxidil, USP 2% w/v Hair regrowth treatment

Use to regrow hair on the scalp

Warnings

For external use only Hammable: Keep away from fire or flame

Do not use if

- your degree of hair loss is different than that shown on the side of this carton, because this product may not work for you
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- · your hair loss is associated with childbirth
- · you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- vou use other medicines on the scalp

Ask a doctor before use if you have heart disease

When using this product

- . do not apply on other parts of the body
- · avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. You may need to use this product 2 times a day for at least 4 months before you see results.
- the amount of hair regrowth is different for each person. This product will not work for everyone.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- · scalp irritation or redness occurs
- unwanted facial hair growth occurs
- · you do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply one mL with dropper 2 times a day directly onto the scalp in the hair loss area
- using more or more often will not improve results
- · continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- in clinical studies of mostly white women aged 18 to 45 years with mild to moderate degree
 of hair loss, the following response to 2% minoxidil topical solution was reported: 19% of women reported moderate hair regrowth after using 2% minoxidil topical solution for 8 months (19% had moderate regrowth; 40% had minimal regrowth). This compares with 7% of women reporting moderate hair regrowth after using the placebo, the liquid without minoxidil in it, for 8 months (7% had moderate regrowth; 33% had minimal regrowth).
- store at controlled room temperature 20° to 25°C (68° to 77°F).

Drug Facts (continued)

Inactive ingredients alcohol (60% v/v), propylene glycol, purified water

Questions?

1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

Minoxidil Topical Solution is for women who have a general thinning of hair on the top of the scalp as shown.

Minoxidil Topical Solution has been shown to regrow hair in women with the following degrees of thinning







If women have more hair loss than shown above, Minoxidil Topical Solution may not work.



Do not use if you have patchy hair loss as shown above.

This Package Contains:

- Two 60 mL (2 fl oz) bottles of Minoxidil Topical Solution (two month supply).
- · One child-resistant dropper applicator
- Information leaflet with complete directions on how to use and obtain best results

*This product is not affiliated with, manufactured by, or produced by the makers or owners of Rogaine®

Distributed by: Pharmacy Value Alliance, LLC 407 East Lancaster Avenue, Wayne, PA 19087



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

Made in Canada 00660616 R0417 VC113940

MINOXIDIL FOR WOMEN

minoxidil solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-006
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)	MINOXIDIL	2 g in 100 mL

Inactive Ingredients		
Ingredient Name	Strength	
ALCOHOL (UNII: 3K9958V90M)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:68016- 006-64	2 in 1 CARTON	01/01/2000		
1		60 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074588	01/01/2000	

Revised: 11/2022 Chain Drug Consortium, LLC