

HALORINNA SUN-SCREEN TRAVEL SPF50- sun screen lotion
Guangdong Bisutang Biotechnology Co., LTD

Directions

- Apply generously and evenly to face and neck 15-20 minutes before sun exposure.
- Reapply at least every 2 hours
- Use a water-resistant sunscreen when swimming or sweating.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early aging. To reduce this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limiting time in the sun, especially from 10 a.m. - 2 p.m.
 - Wear long-sleeved shirts, pants, hats, and sunglasses.
- Children under 6 months: Ask a doctor.

Uses

Helps prevent sunburn

If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early aging caused by the sun.

WATER
GLYCERIN
CYCLOPENTASILOXANE
CETEARYL ALCOHOL
AMMONIUM POLYACRYLOYLDIMETHYL TAURATE
PEG-100 STEARATE
GLYCERYL STEARATE
STEARIC ACID
CYCLOHEXASILOXANE
PHENOXYETHANOL
HYDROXYACETOPHENONE
POTASSIUM CETYL PHOSPHATE
SODIUM HYDROXIDE
FRAGRANCE
ALLANTOIN
SILICA
XANTHAN GUM
DIMETHICONE
DISODIUM EDTA
BHT
DIPOTASSIUM PHOSPHATE
SODIUM CARBONATE

Warnings

Do not use on damaged or broken skin.

For external use only.

When using this product keep out of eyes.

If product gets into the eyes, rinse with water to remove

Stop use and consult a doctor if rash occurs.

ETHYLHEXYL METHOXYCINNAMATE-----7.5

OCTOCRYLENE -----3.4937

PHENYLBENZIMIDAZOLE SULFONIC ACID----- 2

BUTYL METHOXYDIBENZOYLMETHANE -----2

CI 77891 -----1.8

Keep out of reach of children.

Sunscreen



HALORINNA SUN-SCREEN TRAVEL SPF50

sun screen lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84753-040
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLBENZIMIDAZOLE SULFONIC ACID (UNII: 9YQ9DI1W42) (PHENYLBENZIMIDAZOLE SULFONIC ACID - UNII:9YQ9DI1W42)	PHENYLBENZIMIDAZOLE SULFONIC ACID	2 g in 100 mL
ETHYLHEXYL METHOXYCINNAMATE (UNII: 4Y5P7MUD51) (ETHYLHEXYL METHOXYCINNAMATE - UNII:4Y5P7MUD51)	ETHYLHEXYL METHOXYCINNAMATE	7.5 g in 100 mL
BUTYL METHOXYDIBENZOYLMETHANE (UNII: G63QQF2NOX) (BUTYL METHOXYDIBENZOYLMETHANE - UNII:G63QQF2NOX)	BUTYL METHOXYDIBENZOYLMETHANE	2 g in 100 mL
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	3.4937 g in 100 mL
CI 77891 (UNII: 15FIX9V2JP) (CI 77891 - UNII:15FIX9V2JP)	CI 77891	1.8 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOPENTASILOXANE (UNII: 0THT5PCI0R)	
CETEARYL ALCOHOL (UNII: 2DMT128M1S)	
XANTHAN GUM (UNII: TTV12P4NEE)	
ALLANTOIN (UNII: 344S277G0Z)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
POTASSIUM CETYL PHOSPHATE (UNII: 03KCY6P7UT)	
FRAGRANCE FLORAL ORC0902236 (UNII: R66Z4YW3X0)	
BHT (UNII: 1P9D0Z171K)	
GLYCERIN (UNII: PDC6A3C0OX)	
AMMONIUM POLYACRYLOYLDIMETHYL TAURATE (UNII: F01RIY4371)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
GLYCERYL STEARATE (UNII: 230OU9XXE4)	
CYCLOHEXASILOXANE (UNII: XHK3U310BA)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
HYDROXYACETOPHENONE (UNII: G1L3HT4CMH)	
PEG-100 STEARATE (UNII: YD01N1999R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
WATER (UNII: 059QF0KO0R)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TRIAMCINOLONE ACETONIDE 21-DIPOTASSIUM PHOSPHATE (UNII: FP1AQ8LL5O)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84753-040-01	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/23/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	03/23/2026	

Labeler - Guangdong Bisutang Biotechnology Co., LTD (447075429)

Establishment

Name	Address	ID/FEI	Business Operations
Guangdong Bisutang Biotechnology Co., LTD		447075429	manufacture(84753-040)

Revised: 3/2026

Guangdong Bisutang Biotechnology Co., LTD