

**LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride solution**  
**DOLGENCORP INC**

-----  
**Dollar General -Loperamide HCL Oral Solution, USP**

***Drug Facts***

***Active ingredient (in each 7.5 mL)***

Loperamide HCl, USP 1 mg

***Purpose***

Anti-diarrheal

***Use***

controls symptoms of diarrhea, including Travelers' Diarrhea

***Warnings***

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9 years to 11 years (60 lbs to 95 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6 years to 8 years (48 lbs to 59 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2 years to 5 years (34 lbs to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- each 30 mL contains: **sodium 7 mg**
- store at 20°C to 25°C (68°F to 77°F)
- **do not use if outer neck band is broken or missing.**

**Inactive ingredients**

anhydrous citric acid, D&C yellow no. 10, FD&C blue no. 1, flavadew coolmint, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, plain caramel powder, propylene glycol, purified water, simethicone emulsion, sodium benzoate, sucralose, titanium dioxide, xanthan gum

**Questions or comments?**

call toll-free **1-888-235-2466**

**Do not use if outer neck band is broken or missing**

**PEEL OPEN TO READ COMPLETE DIRECTIONS AND WARNINGS BEFORE PURCHASE**

\*This product is not manufactured or distributed by the owners of Imodium<sup>®</sup> A-D.

**DISTRIBUTED BY OLD EAST MAIN CO.**

**100 MISSION RIDGE**

**GOODLETTSVILLE, TN 37072**

**MADE IN INDIA**

**100% Satisfaction Guaranteed!**

(888) 309-9030

984006960

L0001226 R0226 A1791

Code: TN/Drugs/TN00002222/2006

**PRINCIPAL DISPLAY PANEL-4 fl oz**

**DG™ health**

**\*Compare to the active ingredient of Imodium® A-D**

**children's**

**For Ages 6**

**Years and Up**

**Anti-Diarrheal**


**Loperamide Hydrochloride**

**Oral Solution, USP 1 mg per 7.5 mL**

**Controls the symptoms of diarrhea**

**Mint Flavor**

4 fl oz (120 mL)



**Do not use if outer neck band is broken or missing**

**DG™ health**

**\*Compare to the active ingredient of Imodium® A-D**

**Children's** For Ages 6 Years and Up  
**Anti-Diarrheal**

Loperamide Hydrochloride  
Oral Solution, USP 1 mg per 7.5 mL

**Controls the symptoms of diarrhea**



Mint Flavor

**4 FL OZ (120 mL)**

**Drug Facts**

Active ingredient (in each 7.5 mL)	Purpose
Loperamide HCl, USP 1 mg.....	Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  
**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use** if you have bloody or black stool

\*This product is not manufactured or distributed by the owners of Imodium® A-D.  
DISTRIBUTED BY OLD EAST MAIN CO. 948006960.L0001226.R0226.A1791  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072  
MADE IN INDIA



**100% Satisfaction Guaranteed!**  
(888) 309-9030

UVZ  
15 mm x 33 mm  
(Lot and Exp online Printing)

PEEL OPEN TO READ COMPLETE DIRECTIONS AND WARNINGS BEFORE PURCHASE

Top Layer Glue Side

<p><b>Drug Facts</b> (continued)</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ fever            ■ mucus in the stool</li> <li>■ a history of liver disease</li> <li>■ a history of abnormal heart rhythm</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b> taking a prescription drug. Loperamide may interact with certain prescription drugs.</p> <p><b>When using this product</b> tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.</p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ symptoms get worse</li> <li>■ diarrhea lasts for more than 2 days</li> <li>■ you get abdominal swelling or bulging. These may be signs of a serious condition. ▶</li> </ul>	<p><b>Drug Facts</b> (continued)</p> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ <b>drink plenty of clear fluids to help prevent dehydration caused by diarrhea</b></li> <li>■ find right dose on chart. If possible, use weight to dose; otherwise use age.</li> <li>■ shake well before using</li> <li>■ use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.</li> <li>■ mL = milliliter ▶</li> </ul>
--	--

No Text area

12 mm

## Base Layer

<p style="writing-mode: vertical-rl; transform: rotate(180deg);">No Text area</p> <p style="writing-mode: vertical-rl; transform: rotate(180deg);">T00319R0925</p>	<p><b>Drug Facts</b> (continued)</p> <table border="1" style="width: 100%;"> <tr> <td style="width: 30%;">adults and children 12 years and over</td> <td>30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours</td> </tr> <tr> <td>children 9 years to 11 years (60 lbs to 95 lbs)</td> <td>15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours</td> </tr> <tr> <td>children 6 years to 8 years (48 lbs to 59 lbs)</td> <td>15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours</td> </tr> <tr> <td>children 2 years to 5 years (34 lbs to 47 lbs)</td> <td>ask a doctor ▶</td> </tr> </table>	adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours	children 9 years to 11 years (60 lbs to 95 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours	children 6 years to 8 years (48 lbs to 59 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours	children 2 years to 5 years (34 lbs to 47 lbs)	ask a doctor ▶	<p><b>Drug Facts</b> (continued)</p> <table border="1" style="width: 100%;"> <tr> <td style="width: 30%;">children under 2 years (up to 33 lbs)</td> <td>do not use</td> </tr> </table> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ each 30 mL contains: <b>sodium 7 mg</b></li> <li>■ store at 20°C to 25°C (68°F to 77°F)</li> <li>■ <b>do not use if outer neck band is broken or missing.</b></li> </ul> <p><b>Inactive ingredients</b> anhydrous citric acid, D&amp;C yellow no. 10, FD&amp;C blue no. 1, flavadew coolmint, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, plain caramel powder, propylene glycol, purified water, simethicone emulsion, sodium benzoate, sucralose, titanium dioxide, xanthan gum</p> <p><b>Questions or comments?</b> call toll-free 1-888-235-2466</p>	children under 2 years (up to 33 lbs)	do not use
	adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours										
	children 9 years to 11 years (60 lbs to 95 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours										
	children 6 years to 8 years (48 lbs to 59 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours										
	children 2 years to 5 years (34 lbs to 47 lbs)	ask a doctor ▶										
children under 2 years (up to 33 lbs)	do not use											

12 mm

## LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-155
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
CARMEL (UNII: T9D99G2B1R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	

**Product Characteristics**

Color		Score	
Shape		Size	
Flavor	MINT (flavadev coolmint)	Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-155-55	120 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/30/2026	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA220100	03/30/2026	

**Labeler** - DOLGENCORP INC (068331990)**Registrant** - Bionpharma Inc. (079637826)

---

## Establishment

Name	Address	ID/FEI	Business Operations
OrBion Pharmaceuticals Private Limited		854403569	manufacture(55910-155)

Revised: 3/2026

DOLGENCORP INC