

# **STOMACH RELIEF- bismuth subsalicylate liquid QUALITY CHOICE (Chain Drug Marketing Association)**

-----  
**Drug Facts**

## **Active ingredient (in each 30 mL)**

Bismuth subsalicylate 525 mg

## **Purpose**

Upset stomach reliever/Antidiarrheal

## **Uses**

relieves

- diarrhea
- travelers' diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - fullness
  - belching

## **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

## **Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

## **Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

## **Ask a doctor before use if you have**

- fever
- mucus in the stool

## **Ask a doctor or pharmacist before use if you are**

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

## **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

## **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## **Directions**

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adult and children 12 years and over:
  - 30 mL (1 dose) every 1/2 or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
  - 30 mL (1 dose) every 1/2 hour as needed for overindulgence (unset stomach, heartburn, indigestion, nausea)
  - do not exceed 8 doses (240 mL) in 24 hour
  - use until diarrhea stop but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration cause by diarrhea

## **Other information**

- **each 30 mL contains:** sodium 11 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

## **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

## **Questions or comments?**

Call **1-800-935-2362** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

**\*Compare to the Active Ingredient in Pepto-Bismal®**

### **Regular Strength**

#### **Stomach Relief**

Stomach Relief

Bismuth Subsalicylate | Stomach Reliever,

Antidiarrheal

For Relief of:

heartburn

Indigestion

Nausea

Upset Stomach

Diarrhea

Alcohol Free

Sugar Free

Cherry Flavor

FL OZ (mL)

\*This product is not manufactured or distributed by Procter & Gamble Company. Pepto-Bismol® is a registered trademark of th Procter & Gamble Company

**TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

Distributed by C.D.M.A., Inc©

43157 W 9 Mile Rd

Novi, MI 48375

[www.qualitychoice.com](http://www.qualitychoice.com)

# Package Label

**Drug Facts** (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ mL = milliliter ■ shake well before using  
 ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years of age: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL contains: sodium 11 mg ■ each 30 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

**Questions or comments?**  
 Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto Bismol® is a registered trademark of The Procter & Gamble Company.

PLD-A493A  
LB008418



NDC 63868-832-08

\*Compare to the Active Ingredient in Pepto Bismol®

## Regular Strength Stomach Relief

### Stomach Relief

Bismuth Subsalicylate | Upset Stomach Reliever, Antidiarrheal

**For Relief of:**  
 Heartburn  
 Indigestion  
 Nausea  
 Upset Stomach  
 Diarrhea

Alcohol Free  
 Sugar Free

Cherry Flavor

8 fl oz (237 mL)

**Drug Facts** TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

**Active ingredient (in each 30 mL dose)** **Purpose**  
 Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal

**Uses** relieves ■ diarrhea ■ travelers' diarrhea  
 ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ gas ■ indigestion ■ nausea ■ fullness ■ belching

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert: Contains salicylate. Do not take if you are:**  
 ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use** if you have  
 ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have**  
 ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days



6 35515 95562 2

QUALITY CHOICE Stomach Relief

## STOMACH RELIEF

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-832
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-832-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/26/2021	03/26/2026

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/26/2021	03/26/2026

**Labeler** - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 10/2024

QUALITY CHOICE (Chain Drug Marketing Association)