

**DR. SIMI STOMACH RELIEF- bismuth subsalicylate 525 mg suspension**  
**Farmacias De Similares TX LLC**

-----  
**Dr. Simi Stomach Relief**

***Drug Facts***

***Active ingredient***  
***(in each 30 mL)***

Bismuth subsalicylate 525 mg

***Purpose***

Upset stomach reliever and antidiarrheal

***Uses***

relieves: ■ diarrhea ■ heart burn ■ indigestion  
■ nausea ■ upset stomach associated with these symptoms

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- black or bloody stool

**Ask a doctor before use if you have**

- fever ■ mucus in stool

**Ask a doctor or pharmacist before use if you are taking any drug for**

- anticoagulation (thinning of the blood)
- diabetes ■ gout ■ arthritis

**When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- diarrhea lasts more than 2 days
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs

**If pregnant or breast feeding,**

ask health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- shake well before use
- use dose cup or tablespoon (TBSP)
- adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1/2 to 1 hour as needed
- do not exceed 8 doses (240 mL or 16 TBSP) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

***Other information***

- **each 30 mL dose cup contains:** sodium 13 mg, salicylate 256 mg
- protect from freezing avoid excessive heat (over 104°F or 40C)
- low sodium

***Inactive ingredients***

benzoic acid, D&C red #22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

***Questions or comments?***

(855)-280-1079

# Principal Display Panel

**NDC 85786-802-12**

DR. SIMI

Regular Strength

Stomach Relief

Bismuth Subsalicylate 525 mg

Antidiarrheal /Upset Stomach Reliever

Relieves:

- Upset Stomach ■ Nausea ■ Heartburn
- Indigestion ■ Diarrhea

12 fl oz (354 mL)

**TAMPER EVIDENT:** Do not use if imprinted shrinkband is missing or broken

\*Compare to the active ingredient in PEPTO-BISMOL™

\*This product is not manufactured or distributed by Procter&Gamble the distributor of Pepto-Bismol®

Distributed by:

FARMACIAS DE SIMILARES TX LLC.

800 Brazos St Ste 260, Austin, TX, 78701, USA

customerservice@drsimi.com

www.drsimi.com

## Immediate container labeling

**TAMPER EVIDENT:** Do not use if imprinted shrinkband is missing or broken

NDC 85786-802-12

**DR. SIMI** 

\*Compare to the active ingredient in PEPTO-BISMOL™

**Regular Strength  
Stomach Relief  
Bismuth Subsalicylate 525 mg**

**Antidiarrheal /  
Upset Stomach Reliever**

**Relieves:**

- Upset Stomach ■ Nausea ■ Heartburn
- Indigestion ■ Diarrhea

**12 fl oz  
(354 mL)**

**Drug Facts (continued)**

If pregnant or breast feeding, ask health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ Shake well before use  
■ use dose cup or tablespoon (TBSP)  
■ adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1/2 to 1 hour as needed  
■ do not exceed 8 doses (240 mL or 16 TBSP) in 24 hours  
■ use until diarrhea stops but not more than 2 days  
■ children under 12 years: ask a doctor  
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**  
■ each 30 mL dose cup contains: sodium 13 mg, salicylate 256 mg  
■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)  
■ low sodium

**Inactive ingredients** benzoic acid, D&C red #22, D&C red #28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

**Questions or comments?** (855)-280-1079

\* This product is not manufactured or distributed by Procter&Gamble the distributor of Pepto-Bismol™

LR-233

Distributed by:  
**FARMACIAS DE SIMILARES TX LLC.**  
800 Brazos St Ste 260, Austin, TX, 78701, USA  
customerservice@drsimi.com  
www.drsimi.com

**Drug Facts**

**Active ingredient (in each 30 mL)**  
Bismuth subsalicylate 525 mg.....Upset stomach reliever and antidiarrheal

**Purposes**

**Uses** relieves: ■ diarrhea ■ heart burn ■ indigestion  
■ nausea ■ upset stomach associated with these symptoms

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Contains salicylate. Do not take if you are  
■ allergic to salicylates (including aspirin)  
■ taking other salicylate products  
**Do not use if you have** ■ an ulcer  
■ a bleeding problem ■ black or bloody stool  
**Ask a doctor before use if you have**  
■ fever ■ mucus in stool  
**Ask a doctor or pharmacist before use if you are**  
taking any drug for ■ anticoagulation (thinning of the blood)  
■ diabetes ■ gout ■ arthritis  
**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur  
**Stop use and ask a doctor if**  
■ diarrhea lasts more than 2 days  
■ symptoms get worse or last more than 2 days  
■ ringing in the ears or loss of hearing occurs

# DR. SIMI STOMACH RELIEF

bismuth subsalicylate 525 mg suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:85786-802
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM ANHYDROUS</b> (UNII: I4807BK602)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>SODIUM SALICYLATE</b> (UNII: WQ1H85SYP)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85786-802-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/24/2026	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M008	03/24/2026	

**Labeler** - Farmacias De Similares TX LLC (119489934)

## Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma, Inc.		790523323	manufacture(85786-802)

Revised: 3/2026

Farmacias De Similares TX LLC