

**FAMILY WELLNESS EFFERVESCENT COLD RELIEF ORANGE- aspirin,  
chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent  
Family Dollar Services, Inc.**

-----  
**Family Wellness Effervescent Cold Relief Tablets Orange Flavor 20 Ct.**

**Active ingredients**

**(in each effervescent tablet)**

Aspirin 325 mg **(NSAID)\***

Chlorpheniramine maleate 2 mg

Phenylephrine bitartrate 7.8 mg

\*Nonsteroidal anti-inflammatory drug

**Purpose**

Aspirin 325 mg **(NSAID)\***. . . . .Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg . . . . .Antihistamine

Phenylephrine bitartrate 7.8 mg . . . . .Nasal decongestant

\*Nonsteroidal anti-inflammatory drug

**Uses**

temporarily relieves these symptoms of the common cold:

■ minor aches and pains ■ headache ■ runny nose ■ fever ■ nasal and sinus  
congestion ■ sneezing ■ sore throat

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if**

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

### **Ask a doctor before use if you have**

**Ask a doctor before use if you have** ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

**When using this product** ■ do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by a doctor.

- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away.

■ pain or nasal congestion gets worse or lasts more than 7 days. ■ fever gets worse or lasts more than 3 days. ■ new symptoms occur

■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs ■ nervousness, dizziness or sleeplessness occurs

### **If pregnant or breastfeeding**

**If pregnant or breastfeeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

### **Keep out of reach of children**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

■ do not use more than directed (see overdose warning)

■ adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4-6 hours. Do not take more than 8 tablets in 24 hours.

■ ***children under 12 years: ask a doctor***

### **Other information**

■ **each tablet contains:** sodium 464 mg

■ phenylketonurics: contains phenylalanine 9 mg per tablet

■ store at room temperature (59°-86°F)

### **Inactive ingredients**

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

### **Questions or comments?**

**Call 1-888-228-6937**

**PACKAGE LABEL. PRINCIPAL DISPLAY PANEL**



**Drug Facts (continued)**

more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin
- if you are now taking a prescription or over-the-counter (OTC) drug for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if** stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

**Ask a doctor before use if you have** ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood, diabetes, gout, or arthritis)

**When using this product** ■ do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by a doctor.

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have black or bloody stools

■ have stomach pain that does not get better

**Drug Facts**

**Active ingredients**

**(In each effervescent tablet)**

Aspirin 325 mg (NSAID)  
Chlorpheniramine maleate 2 mg  
Phenylephrine bitartrate 7.8 mg  
Nonsteroidal anti-inflammatory drug

**Purpose**

temporarily relieves these symptoms of the common cold:  
■ aches and pains ■ headache ■ runny nose ■ sore throat ■ sneezing ■ sinus congestion

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing aspirin, ibuprofen, naproxen, or others

**Sore throat warning:** If sore throat is severe, persists for product ■ take more or for a longer time than directed

**FAMILY wellness**™

COMPARE TO ALKA-SELTZER PLUS® COLD ORANGE ZEST™

**COLD RELIEF**

Aspirin (NSAID)/Pain Reliever/Fever Reducer  
Chlorpheniramine Maleate/Antihistamine Phenylephrine Bitartrate/Nasal Decongestant

**For Relief of:**  
Nasal Congestion • Runny Nose  
Headache & Body Ache  
Sore Throat • Sinus Pressure

ORANGE FLAVOR

20 EFFERVESCENT TABLETS

**FAMILY wellness**™

COMPARE TO ALKA-SELTZER PLUS® COLD ORANGE ZEST™

**COLD RELIEF**

Aspirin (NSAID)/Pain Reliever/Fever Reducer  
Chlorpheniramine Maleate/Antihistamine Phenylephrine Bitartrate/Nasal Decongestant

**For Relief of:**  
Nasal Congestion  
Runny Nose • Headache & Body Ache • Sore Throat  
Sinus Pressure

ORANGE FLAVOR

20 EFFERVESCENT TABLETS

OTC network

NDC 55319-875-20

**Drug Facts (continued)**

**Other information**

- each tablet contains: sodium 464 mg
- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

**Inactive ingredients**

acesulfame potassium, aspartame, citric acid, docusate sodium, FD&C red #40, FD&C Yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Questions or comments?**  
Call 1-888-228-6937

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

DISTRIBUTED BY: MIDWOOD BRANDS, LLC  
500 VOLVO PARKWAY, CHESAPEAKE, VA 23320 USA

MANUFACTURED IN THE USA WITH DOMESTIC AND IMPORTED INGREDIENTS.

**NOT 100% SATISFIED?** Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

\*This product is not manufactured or distributed by Bayer, owner of the registered trademark Alka-Seltzer Plus® Cold Orange Zest.

SKU 0900495

PN: 87560 FA C2  
R: 06/19


**Drug Facts (continued)**

- an allergic reaction occurs. Seek medical help right away. ■ pain or nasal congestion gets worse or lasts more than 7 days. ■ fever gets worse or lasts more than 3 days. ■ new symptoms occur ■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs ■ nervousness, dizziness or sleeplessness occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4-6 hours. Do not take more than 8 tablets in 24 hours.
- children under 12 years: ask a doctor



# FAMILY WELLNESS EFFERVESCENT COLD RELIEF ORANGE

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-875
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>PHENYLEPHRINE BITARTRATE</b> (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

## Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	ORANGE	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-875-20	10 in 1 CARTON	09/15/2011	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/15/2011	

**Labeler** - Family Dollar Services, Inc. (024472631)

Revised: 12/2023

Family Dollar Services, Inc.