

**FAMILY WELLNESS EFFERVESCENT COLD RELIEF ORANGE- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent Family Dollar Services, Inc.**

-----  
**Family Wellness Effervescent Cold Relief Tablets Orange Flavor 20 Ct.**

**Active ingredients**

**(in each effervescent tablet)**

Aspirin 325 mg **(NSAID)\***

Chlorpheniramine maleate 2 mg

Phenylephrine bitartrate 7.8 mg

\*Nonsteroidal anti-inflammatory drug

**Purpose**

Aspirin 325 mg **(NSAID)\***. . . . .Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg . . . . .Antihistamine

Phenylephrine bitartrate 7.8 mg . . . . .Nasal decongestant

\*Nonsteroidal anti-inflammatory drug

**Uses**

■ temporarily relieves these symptoms due to the common cold:

■ sneezing ■ nasal congestion ■ sore throat ■ headache ■ minor aches and pains ■  
runny nose ■ sinus congestion and pressure

■ temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using

this product ■ take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients.

### **Ask a doctor before use if**

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

### **Ask a doctor before use if you have**

**Ask a doctor before use if you have** ■ asthma ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

#### **When using this product ■ do not exceed recommended dosage**

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away
- pain or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur
- redness or swelling is present ■ ringing in the ears or a loss of hearing occurs ■ nervousness, dizziness or sleeplessness occurs

### **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours
- do not take more than 8 tablets in 24 hours
- ***children under 12 years: ask a doctor***

### **Other information**

**each tablet contains:** sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

### **Inactive ingredients**

acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Questions or comments?

Call 1-888-228-6937

PACKAGE LABEL. PRINCIPAL DISPLAY PANEL

Barcode

**Drug Facts**

**Active ingredients (in each effervescent tablet)**

**Purpose**

**Uses**

**Warnings**

**Drug Facts (continued)**

FAMILY Wellness

EFFERVESCENT

**COLD RELIEF**

Aspirin (NSAID)/Pain Reliever-Fever Reducer  
Chlorpheniramine Maleate/Antihistamine  
Phenylephrine Bitartrate/Nasal Decongestant

**Drug Facts (continued)**

an allergic reaction occurs. Seek medical help right away.

pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

do not use more than directed (see overdose warning)

adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours

do not take more than 8 tablets in 24 hours

children under 12 years: ask a doctor

FAMILY Wellness

COMPARE TO THE ACTIVE INGREDIENTS IN ALKA-SELTZER PLUS® COLD ORANGE ZEST

EFFERVESCENT

**COLD RELIEF**

Aspirin (NSAID)/Pain Reliever-Fever Reducer  
Chlorpheniramine Maleate/Antihistamine  
Phenylephrine Bitartrate/Nasal Decongestant

**RELIEVES:**

- Nasal congestion
- Runny nose
- Sneezing
- Headache
- Sinus pain & pressure

**ORANGE FLAVOR**

20 EFFERVESCENT TABLETS

NDC 55319-893-30

**Drug Facts (continued)**

**Other information**

each tablet contains: sodium 464 mg

phenylephrine contains phenylephrine 9 mg per tablet

store at room temperature (69°-86°F)

**Inactive ingredients**

acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Questions or comments?**

Call 1-888-228-6937

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

DISTRIBUTED BY: MIDWOOD BRANDS, LLC  
300 VOLVO DRIVE, CHESAPEAKE, VA 23020 USA

MANUFACTURED IN THE USA WITH DOMESTIC AND IMPORTED INGREDIENTS

\*This product is not manufactured, distributed or labeled by the maker of Alka-Seltzer Plus® Cold Orange Zest.

NOT 100% SATISFIED? Return within 30 days to the store of purchase for a refund (with receipt) or exchange.

SKU 937866

PN: 87560 FD C1  
REV: 05/23

Barcode

# FAMILY WELLNESS EFFERVESCENT COLD RELIEF ORANGE

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55319-875
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>PHENYLEPHRINE BITARTRATE</b> (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-875-20	10 in 1 CARTON	10/19/2023	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	10/19/2023	

**Labeler** - Family Dollar Services, Inc. (024472631)

Revised: 7/2024

Family Dollar Services, Inc.