#### PURELIFE TOPICAL ANESTHETIC- benzocaine gel PureLife Dental

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### **Active Ingredients**

Benzocaine 200mg (in each g)

## **Purpose**

Oral Anesthetic

#### Use

For oral mucosa only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

#### Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

#### Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally

swallowed, get medical help or contact a Poison Control Center right away. **If pregnant or breast feeding,** ask a physician before use.

Keep out of reach of children.

#### **Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

## Other Information

Store at 59° - 86°F (15°-30°C). Protect from freezing.

# **Inactive Ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

#### Questions or comments?

877-777-3303 or purelifedental.com

#### MANUFACTURED FOR PureLife, LLC., Carson, CA 90810 www.PureLifeDental.com 877-777-3303

# **TOPICAL ANESTHETIC GEL** Benzocaine 20%





Gluten Free 1oz (30ml)

mouth symptoms do not improve f swelling, rash or fever develops, redness persists or worsens; or directed by a physician. If sore n 7 days; irritation, pain, or see your physician promptly

in children under 2 years of age contact occurs, flush with water. When using this product Avoid contact with eyes. If for teething

Do not exceed recommended Poison Control Center right away pain is accidentally swallowed, dosage. If more than used for aet medical help or contact a

may cause methemoglobinemia a serious condition that must be

warning: Use of this product

Methemoglobinemia

Warnings

f pregnant or breast feeding, Keep out of reach of children. ask a physician before use.

Directions

attention if you or a child in your

pale, gray or blue colored skin

care develops:

occur even if you have used this

product before. Stop use and

seek immediate medical

reduces the amount of oxygen

treated promptly because it

carried in the blood. This can

 Apply only amount needed to children under 2 years of age: the oral mucosa to prevent or relieve pain.

store at 59° - 86° F (15 - 30° C). Other information do not use

dizziness or lightheadedness

 shortness of breath rapid heart rate

headache

cyanosis)

Allergy Alert: Do not use on

fatigue or lack of energy

patients with a history of

allergies to local anesthetics

such as procaine, butacaine,

nactive ingredients Protect from freezing.

tartrazine), yellow #6, as a color blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, sodium saccharin. May contain lavoring, PEG 3350, PEG 400,

Questions or comments? additive.

877-777-3303 or purelifedental.com

99192007 Rev 10/2018

# benzocaine or other "caine"

anesthetics.

 for more than 7 days unless Do not use

# PURELIFE TOPICAL ANESTHETIC

temporary relief of pain due

to minor dental procedures

benzocaine gel

Benzocaine 200 mg *(in each g)* 

Active ingredients

**Orug Facts** 

**Product Information** 

HUMAN OTC DRUG NDC:68987-001 **Product Type** Item Code (Source)

**Route of Administration** DENTAL

For oral mucosal use only, as

Oral Anesthetic

Purpose

directed by dentist. For the

**Active Ingredient/Active Moiety** 

**Basis of Strength Ingredient Name** Strength

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) **BENZ OCAINE** 200 mg in 1 g

**Inactive Ingredients** 

**Ingredient Name** Strength POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) SACCHARIN SODIUM (UNII: SB8ZUX40TY)

FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	STRAWBERRY	Imprint Code	
Contains			

	P	Packaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
l	1	NDC:68987-001- 30	30 g in 1 JAR; Type 0: Not a Combination Product	10/01/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	07/01/2013	

# Labeler - PureLife Dental (828690904)

Revised: 1/2022 PureLife Dental