

WISPIRIN- aspirin tablet
Wisdrug AG

Aspirin 81 mg (NSAID)*

Pain reliever / fever reducer

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if the stomach bleeding warning applies to you.

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- anticoagulation (thinning of the blood)
- gout
- diabetes
- arthritis

Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right

away.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

You may report side effects to FDA at 1-800-FDA-1088.

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

- drink a full glass of water with each dose
- before use, disperse the tablet in water immediately and take the suspension promptly, or swallow the tablet whole with water
- do not chew, crush, or split
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours while symptoms persist; do not exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: ask a doctor
- store at 20-25°C (68-77°F)
- protect from moisture
- do not use if imprinted safety seal under cap is broken or missing

Hypromellose, Talc, Hypromellose acetate succinate, Hypromellose phthalate, Triethyl citrate, Microcrystalline cellulose, Povidone K30, Crospovidone, Stearic acid, Aspartame, Anhydrous citric acid, Copovidone, Polysorbate 80, Methacrylic acid - ethyl acrylate copolymer, Glyceryl monostearate.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

- temporarily relieves minor aches and pains
- reduces fever
- for adults who have had a heart attack or stroke: helps prevent another heart attack or stroke (under a doctor's supervision)

Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. Ask your doctor about other uses for this product.

wispirin™
Low Dose Aspirin

ASPIRIN

DELAYED-RELEASE
Tablets for Suspension

81 mg

Pain reliever / fever reducer
Delayed action. Not for fast relief of headache or other symptoms needing immediate relief.

Net contents: 100 tablets

Talk to your doctor or other healthcare provider before using this product for your heart.

Drug Facts	
Active ingredient (in each tablet) Aspirin 81 mg (NSAID)*	Purpose Pain reliever / fever reducer
*nonsteroidal anti-inflammatory drug	
Uses	
<ul style="list-style-type: none"> temporarily relieves minor aches and pains reduces fever for adults who have had a heart attack or stroke: helps prevent another heart attack or stroke (under a doctor's supervision) Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. Ask your doctor about other uses for this product.	
Warnings	
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.	
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: <ul style="list-style-type: none"> hives facial swelling asthma (wheezing) shock 	
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take a blood thinner (anticoagulant) or steroid drug 	

Drug Facts (continued)
Warnings (continued) <ul style="list-style-type: none"> take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if the stomach bleeding warning applies to you, <ul style="list-style-type: none"> you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic you have asthma
Ask a doctor or pharmacist before use if you are taking a prescription drug for: <ul style="list-style-type: none"> anticoagulation (thinning of the blood) gout diabetes asthma
Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away.
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> stomach pain black stools bloody or black stools have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days redness or swelling is present new symptoms occur ringing in the ears or loss of hearing occurs You may report side effects to FDA at 1-800-FDA-1088.

Drug Facts (continued)
Warnings (continued) <ul style="list-style-type: none"> If pregnant or breastfeeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> drink a full glass of water with each dose before use, dispense the tablet in water immediately and take the suspension promptly, or swallow the tablet whole with water do not chew, crush, or split adults and children 12 years and over: take 4 to 8 tablets every 4 hours while symptoms persist; do not exceed 48 tablets in 24 hours unless directed by a doctor children under 12 years of age: ask a doctor
Other information
<ul style="list-style-type: none"> store at 20-25°C (68-77°F) protect from moisture do not use if implanted safety seal under cap is broken or missing
Inactive ingredients
Hypromellose, Talc, Hypromellose acetate succinate, Hypromellose phthalate, Triethyl citrate, Microcrystalline cellulose, Povidone K30, Croscopolvidone, Stearic acid, Aspartame, Anhydrous citric acid, Copovidone, Polyethylene 80, Methacrylic acid - ethyl acrylate copolymer, Glycerol monostearate.
Questions or comments?
Call 1-877-258-0839 Manufacturer: Windring AG Address: Landis-Gyr-Strasse 1, Zug, 6300, Switzerland (CHE)

WISPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87580-001
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg in 265.45 mg

Inactive Ingredients

Ingredient Name	Strength
COPOVIDONE K25-31 (UNII: D9C330MD8B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE K30 (UNII: U725QWY32X)	
TALC (UNII: 7SEV7J4R1U)	
CROSPVIDONE (UNII: 2S7830E561)	
HYPROMELLOSE PHTHALATE (31% PHTHALATE, 40 CST) (UNII: G4U024CQK6)	
ASPARTAME (UNII: Z0H242BBR1)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
HYPROMELLOSE ACETATE SUCCINATE 12070923 (3 MM2/S) (UNII: 36BGF0E889)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	

Product Characteristics

Color	white	Score	score with uneven pieces
Shape	ROUND	Size	8mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87580-001-01	15927 mg in 1 BOTTLE; Type 0: Not a Combination Product	04/15/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/15/2026	

Labeler - Wisdrug AG (481044776)

Registrant - Wisdrug AG (481044776)

Establishment

Name	Address	ID/FEI	Business Operations
Wisdrug AG		481044776	manufacture(87580-001)

Revised: 4/2026

Wisdrug AG