

**DAY TIME GELCAP - acetaminophen, dextromethorphan, phenylephrine capsule, liquid filled
Kareway Product, Inc.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each GelCap)

Acetaminophen 325mg

Dextromethorphan HBr 10mg

Phenylephrine HCl 5mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal decongestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Enter section text here

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Sore throat warning:

If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm(mucus)
- Persistent or chronic cough as occurs with smoking, asthma, emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children

Enter section text here

Overdose warning:

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical to adults and children even if you do not notice any signs or symptoms.

Directions

- take only as directed-see Overdose warning
- do not exceed 4 doses per 24 hours

| | |
|-------------------------------------|----------------------------------|
| adults and children 12 yrs and over | 2 GelCaps with water every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- store at room temperature

Inactive ingredients

FDC red no. 40, FDC yellow no. 5, dioxide, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water and sobital special

Daytime gelcap

Pure-Aid™
GelCap

Day Time

NEW FORMULA WITH PHENYLEPHRINE

Multi-Symptom
COLD/FLU Relief

Pure-Aid™
GelCap

Multi-Symptom COLD/FLU Relief
Day Time
NEW FORMULA WITH PHENYLEPHRINE

Pure-Aid™

NON-DROWSY

Day Time

NEW FORMULA WITH PHENYLEPHRINE

GelCap

See new warnings and dosing information

*Compare to
active ingredients
in DayQUIL®

Relieves

- Fever
- Aching
- Stuffy Head
- Coughing
- Sore Throat

Without Drowsiness!

Acetaminophen

Aches/Fever reducer

Dextromethorphan Hbr

Cough suppressant

Phenylephrine

Nasal congestion

Multi-Symptom
COLD/FLU Relief



10 GELCAPS

Pure-Aid™
GelCap

Day Time

TAMPER EVIDENT: This package is safety sealed and child resistant. Use only if blisters are intact. If difficult to open, use scissors.



| | |
|--|---|
| <p>Drug Facts</p> <p>Failure to follow these warnings could result in serious consequences.</p> <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless (children) or symptoms get worse or last more than 5 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition ■ If pregnant or breast-feeding, ask a health professional before use. ■ Keep out of reach of children. Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.</p> <p>Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to enlarged prostate gland ■ cough ■ Persistent or chronic cough as occurs with smoking, asthma, or emphysema ■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> | |
| <p>Active ingredients (in each GelCap)</p> <p>Acetaminophen 325 mg Dextromethorphan Hbr 10 mg Phenylephrine HCl 5 mg Nasal decongestant</p> | <p>Warnings Liver warning: this product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily amount for this product ■ Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to enlarged prostate gland ■ cough ■ Persistent or chronic cough as occurs with smoking, asthma, or emphysema ■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> |
| <p>Drug Facts (continued)</p> <p>Purpose</p> <p>Relieves fever, aches, and pain. ■ Relieves cold and flu symptoms, including cough, sore throat, and nasal congestion.</p> | <p>Directions ■ Take only as directed—see Overdose warning ■ Do not exceed 4 doses per 24 hours ■ Do not exceed 4 doses per 24 hours ■ adults & children 12 yrs & over: 2 GelCaps with water every 6 hrs ■ children 4 to under 12 yrs: ask a doctor ■ children under 4 yrs: do not use</p> <p>Other information ■ store at room temperature ■ Do not use ■ ask a doctor ■ children under 4 yrs: do not use</p> <p>Inactive ingredients FD&C red no. 40, FD&C yellow no. 5, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, and sorbitol special</p> <p><small>*This product is not manufactured by Procter & Gamble, owner of the registered trademark Dayquil®.</small></p> <p>Exclusively distributed by: KARVE WAVE Karvey Product Inc., Compton, CA 90220 Made in Korea</p> |

Multi-Symptom COLD/FLU Relief
Day Time
NEW FORMULA WITH PHENYLEPHRINE

Pure-Aid™
GelCap

DAY TIME GELCAP

acetaminophen, dextromethorphan, phenylephrine capsule, liquid filled

Product Information

| | | | |
|-------------------------|----------------|--------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:67510-0523 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| FD&C YELLOW NO. 5 (UNII: I753WB2F1M) | |
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A) | |
| POVIDONE (UNII: FZ989GH94E) | |
| PROPYLENE GLYCOL, (R)- (UNII: 602HN5L69H) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITOL (UNII: 506T60A25R) | |

Product Characteristics

| | | | |
|----------|---------|--------------|----------|
| Color | orange | Score | no score |
| Shape | CAPSULE | Size | 24mm |
| Flavor | | Imprint Code | DYTIME |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|----------------------|----------------------|--------------------|
| 1 | NDC:67510-0523-1 | 1 in 1 BOX | | |
| 1 | | 10 in 1 BLISTER PACK | | |
| 2 | NDC:67510-0523-2 | 1 in 1 BOX | | |
| 2 | | 12 in 1 BLISTER PACK | | |
| 3 | NDC:67510-0523-0 | 2 in 1 BOX | | |
| 3 | | 20 in 1 BLISTER PACK | | |
| 4 | NDC:67510-0523-4 | 2 in 1 BOX | | |
| 4 | | 24 in 1 BLISTER PACK | | |

Marketing Information

| | | | |
|--------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|

Labeler - Kareway Product, Inc. (121840057)

Revised: 5/2011

Kareway Product, Inc.