

**SODIUM CHLORIDE OPHTHALMIC 5 PERCENT HYPERTONICITY EYE- sodium chloride ointment**

**Hanlim**

-----  
**Hanlim (MFG) for Walgreens Hypertonicity (PLD)**

**Walgreens Sodium Chloride Ophthalmic**

**Ointment 5% Hypertonicity Eye Ointment 3.5g**

**NDC 0363-7500-50**

***Drug Facts***

***Active ingredient***

Sodium Chloride, 50 mg (5%)

***Purpose***

Hypertonicity Agent

***Use***

- For the temporary relief of corneal edema.

***Warnings***

- **For use in the eyes only.**
- Retain outer carton for full product drug facts.

***Do not use***

this product except under the advice and supervision of a doctor.

***When using this product***

- avoid contamination, do not touch tip of container to any surface.
- replace cap after use.
- this product may cause temporary burning and irritation on being instilled into the eye.

***Stop use and ask a doctor if***

- you experience eye pain, changes in vision, continued redness or irritation of the eye.
- condition worsens or persists for more than 72 hours.

**KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- pull down lower lid of the affected eye(s) and apply a small amount (1/4 inch) to the inside of the eyelid and apply every 3-4 hours, or as directed by a doctor.

***Other information***

- Store at room temperature 15°- 30°C (59°- 86°F).
- Do not freeze.
- Keep tightly closed.
- See crimp of tube or box for lot number and expiration date.

***Inactive ingredients***

lanolin alcohol, mineral oil, water for injection and white petrolatum.

***Questions or comments?***

- 1- 800-925-4733

**NDC 0363-7500-50**

**Sodium Chloride**

**Ophthalmic**

**Ointment, 5%**

**Hypertonicity**

**Eye Ointment**

**NET WT 0.125 OZ (3.5 g)**

Walgreens

NDC 0363-7500-50

Walgreens


Compare to the active ingredient in Muro 128®  
Hypertonicity Ophthalmic Ointment, 5%<sup>††</sup>

# Sodium Chloride Ophthalmic Ointment, 5% Hypertonicity Eye Ointment

- For temporary relief of corneal edema
- Sterile

DISTRIBUTED BY:  
**WALGREEN CO.**  
 200 WILMOT RD.,  
 DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
 walgreens.com  
 ©2022 Walgreen Co.  
**MADE IN SOUTH KOREA**

CE0SC0012WR2



NET WT 0.125 OZ (3.5 g)

LOT & EXP

LEAVE AREA BLANK.  
ALLOCATED FOR  
LOT & EXP CODE

W3ORG0223-F  
REV-0323

### Drug Facts (continued)

#### Other information

- store at 15-25°C (59-77°F)
- DO NOT FREEZE
- keep tightly closed
- do not use if difficult to dispense or visible particles are seen in the product
- do not roll, bend, twist, or fold tube during use

#### Inactive ingredients

lanolin, light mineral oil, water for injection, white petrolatum

#### Questions or comments?

Call 1-888-527-4276

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

<sup>††</sup>This product is not manufactured or distributed by Bausch & Lomb, owner of the registered trademark Muro 128®.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPEN OR NECKBAND ON BOTTOM OF CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**DO NOT USE IF BOTTOM RIDGE OF TUBE IS EXPOSED, OR IF CAP OR TUBE IS DAMAGED.**

### Drug Facts

#### Active ingredient Purpose

Sodium chloride 5%.....Hypertonicity agent

#### Uses

- temporary relief of corneal edema

#### Warnings

For external use only

##### Do not use

- except under the advice and supervision of a doctor

##### When using this product

- it may cause temporary burning and irritation
- replace cap after use
- to avoid contamination do not touch tip of container to any surface

##### Stop use and ask a doctor if

- you experience eye pain, changes in vision, continued redness or irritation of the eye
- condition worsens or persists for more than 72 hours


##### Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions

- pull down the lower lid of the affected eye
- apply a small amount (1/4 inch) of ointment to the inside of eyelid
- apply every 3 or 4 hours or as directed by a doctor

## Tube Label



## SODIUM CHLORIDE OPHTHALMIC 5 PERCENT HYPERTONICITY EYE

sodium chloride ointment

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11716-0750
<b>Route of Administration</b>	OPHTHALMIC		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W471Q8X) (CHLORIDE ION - UNII:Q32Z N48698)	SODIUM CHLORIDE	50 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>LANOLIN ALCOHOLS</b> (UNII: 884C3FA9HE)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11716-0750-0	1 in 1 CARTON	04/04/2026	
1		3.5 g in 1 TUBE; Type 0: Not a Combination		

Product			
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	04/04/2026	

**Labeler** - Hanlim (687986034)

Revised: 4/2026

Hanlim